

For Research Use Only

Beta Catenin Polyclonal antibody

Catalog Number: 51067-2-AP

Featured Product

885 Publications



Basic Information

Catalog Number:

51067-2-AP

Size:

150ul, Concentration: 500 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC058926

GeneID (NCBI):

1499

ENSEMBL Gene ID:

ENSG00000168036

UNIPROT ID:

P35222

Full Name:

catenin (cadherin-associated protein), beta 1, 88kDa

Calculated MW:

781 aa, 86 kDa

Observed MW:

92 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:1000-1:4000

IF-P 1:500-1:2000

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, FC (Intra), IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, ChIP

Species Specificity:

human, mouse, rat, pig

Cited Species:

human, mouse, rat, pig, canine, chicken, bovine, hamster, goat, sheep

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, HeLa cells, HEK-293T cells, NIH/3T3 cells, C6 cells, mouse liver tissue, rat liver tissue

IP: mouse liver tissue,

IHC: mouse liver tissue, human colon cancer tissue, human liver cancer tissue, mouse colon tissue

IF-P: mouse testis tissue, mouse epididymis tissue

IF/ICC: MCF-7 cells, HepG2 cells, HeLa cells, T-47D cells, mouse testis tissue

Background Information

β -Catenin, also known as CTNNB1, is an evolutionarily conserved, multifunctional intracellular protein. β -Catenin is a 92-kDa protein, originally identified in cell adherens junctions (AJs) where it functions to bridge the cytoplasmic domain of cadherins to α -catenin and the actin cytoskeleton. Besides its essential role in the AJs, β -catenin is also a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The Wnt/ β -catenin pathway is also involved in the activation of other intracellular messengers such as calcium fluxes, JNK, and SRC kinases. Deregulation of β -catenin activity is associated with multiple diseases including cancers. (PMID: 22617422; 18334222)

Notable Publications

Author	Pubmed ID	Journal	Application
Min Zhou	36171592	J Exp Clin Cancer Res	WB
Wen-Long Lei	34580275	Cell Death Dis	WB,IF
Ke-Xin Wang	34649212	Phytomedicine	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of Beta Catenin antibody (51067-2-AP; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Beta Catenin transfected HeLa cells.




Various lysates were subjected to SDS PAGE followed by western blot with 51067-2-AP (Beta Catenin antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 51067-2-AP (Beta Catenin antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 51067-2-AP (Beta Catenin antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using Beta Catenin antibody (51067-2-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




IP result of anti-Beta Catenin (IP:51067-2-AP, 4ug; Detection:51067-2-AP 1:500) with mouse liver tissue lysate 6400ug.



Immunofluorescent analysis of (4% PFA) fixed mouse testis tissue using Beta Catenin antibody (51067-2-AP) at dilution of 1:1000 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10⁶ MCF-7 cells were intracellularly stained with 0.2 ug Anti-Human Beta Catenin (51067-2-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Beta Catenin antibody (51067-2-AP) at dilution of 1:400 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), Alpha Tubulin antibody (66031-1-Ig, Clone: 1E4C11, red).