For Research Use Only

p38 MAPK Polyclonal antibody

Catalog Number: 51115-1-AP 31 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

51115-1-AP Size:

GeneID (NCBI):

Antigen affinity purification

Recommended Dilutions:

150ul, Concentration: 500 ug/ml by Nanodrop and 260 ug/ml by Bradford $\,$ UNIPROT ID:

BC031574

WB 1:500-1:1000

method using BSA as the standard;

Q16539 Full Name:

Source: Rabbit

mitogen-activated protein kinase 14

Isotype:

Calculated MW:

360 aa, 41 kDa Observed MW:

38 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human, mouse, rat, pig

human, mouse, rat

Cited Species:

Positive Controls:

WB: mouse ovary tissue, NIH/3T3 cells, PC-12 cells,

RAW 264.7 cells

Background Information

MAPK14(mitogen-activated protein kinase 14) is also named as SAPK2A, p38MAPK, CSBP1, RK, p38, EXIP, Mxi2, CSBP2, PRKM14, PRKM15, CSPB1, p38ALPHA and belongs to the MAP kinase subfamily. MAPK14-signaling is a central pathway for the integration of instructive signals in dendritic cells for T(H)17 differentiation and in flam mation (PMID: 22231518). It plays an important role in the regulation of hematopoietic stem cells elf-renewal and the regulation of hematopoietic stem of the regulation of thein vitro and inhibition of MAPK14 activation with a small molecule inhibitor may represent a novel approach to promote ex vivo expansion of hematopoietic stem cell(PMID:21198398). This protein has 4 isoforms produced by alternative splicing. This antibody is specific to MAPK14.

Notable Publications

Author	Pubmed ID	Journal	Application
Yu Zhou	36430479	Int J Mol Sci	WB
Weizong Wang	35821539	Cardiovasc Drugs Ther	WB
Xiaojuan Peng	35866029	Front Bioeng Biotechnol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 51115-1-AP (p38 MAPK antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 51115-1-AP (p38 MAPK antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.