For Research Use Only

USP22 Polyclonal antibody

Catalog Number:55110-1-AP

Featured Product 15 Publications




Basic Information	Catalog Number: 55110-1-AP	GenBank Accession Number: NM_015276		Purification Method: Antigen affinity purification		
	Silo PAP Size: 150ul , Concentration: 750 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GeneID (NCBI): Rec 23326 WB UNIPROT ID: IP 0		Recommended Dilutions: WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate		
					ubiquitin specific peptidase 22 Calculated MW: 60 kDa	
				Observed MW: 58-60 kDa		
		Applications	Tested Applications:	Positive Controls: WB : A549 cells, HepG2 cells IP : HepG2 cells,		
			WB, IP, ELISA			
			Cited Applications: WB, IHC, IF, IP, CoIP			
Species Specificity: human						
	Cited Species: human, mouse, rat					
Background Information	human, mouse, rat USP22, also named as KIAA 1063 and deubiquitinating component of the tu catalyzes the deubiquitination of bot specific gene promoters by activator.	ranscription regul th histones H2A a s such as MYC, wh vation and cell cy	latory histone acety and H2B, thereby action here it is required for	ng as a coactivator. It is recruited to		
<u> </u>	human, mouse, rat USP22, also named as KIAA 1063 and deubiquitinating component of the tr catalyzes the deubiquitination of bot specific gene promoters by activator nuclear receptor-mediated transactiv 58kDa and 60 kDa. The antibody is sp	ranscription regul ch histones H2A a s such as MYC, wh vation and cell cy pecific to USP22.	latory histone acety and H2B, thereby acti here it is required fo ccle progression. USF	lation (HAT) complex SAGA. USP22 ing as a coactivator. It is recruited to r transcription. USP22 is required for 22 has 2 isoforms with molecular mass		
	human, mouse, rat USP22, also named as KIAA1063 and deubiquitinating component of the to catalyzes the deubiquitination of bot specific gene promoters by activator nuclear receptor-mediated transactive 58kDa and 60 kDa. The antibody is sp Author Pu	ranscription regul th histones H2A a s such as MYC, wh ration and cell cy pecific to USP22.	latory histone acety and H2B, thereby acti here it is required fo rcle progression. USF Journal	lation (HAT) complex SAGA. USP22 ng as a coactivator. It is recruited to r transcription. USP22 is required for P22 has 2 isoforms with molecular mass Application		
	human, mouse, rat USP22, also named as KIAA1063 and deubiquitinating component of the tr catalyzes the deubiquitination of bot specific gene promoters by activator nuclear receptor-mediated transactiv 58kDa and 60 kDa. The antibody is sp Author Pu Dianyun Ren 34	ranscription regul ch histones H2A a s such as MYC, wh vation and cell cy pecific to USP22.	latory histone acety and H2B, thereby acti here it is required fo ccle progression. USF	lation (HAT) complex SAGA. USP22 ing as a coactivator. It is recruited to r transcription. USP22 is required for 22 has 2 isoforms with molecular mass		
	human, mouse, rat USP22, also named as KIAA1063 and deubiquitinating component of the tr catalyzes the deubiquitination of bot specific gene promoters by activator nuclear receptor-mediated transactiv 58kDa and 60 kDa. The antibody is sp Author Pu Dianyun Ren 34 Zelong Han 36	ranscription regul th histones H2A a s such as MYC, wh ration and cell cy pecific to USP22. bmed ID 743406	latory histone acety and H2B, thereby acti here it is required fo rcle progression. USF Journal Mol Oncol	lation (HAT) complex SAGA. USP22 ing as a coactivator. It is recruited to r transcription. USP22 is required for 22 has 2 isoforms with molecular mass Application WB,IP		
	human, mouse, rat USP22, also named as KIAA 1063 and deubiquitinating component of the tr catalyzes the deubiquitination of bot specific gene promoters by activator nuclear receptor-mediated transactiv 58kDa and 60 kDa. The antibody is sp Author Pu Dianyun Ren 34 Zelong Han 36	ranscription regul th histones H2A a s such as MYC, wh ration and cell cy pecific to USP22. bmed ID 743406 3333288	latory histone acety and H2B, thereby acti here it is required for ccle progression. USF Journal Mol Oncol Cell Death Discov	lation (HAT) complex SAGA. USP22 ing as a coactivator. It is recruited to r transcription. USP22 is required for 22 has 2 isoforms with molecular mass Application WB,IP WB,IHC,IF,IP		
Background Information Notable Publications	human, mouse, rat USP22, also named as KIAA 1063 and deubiquitinating component of the tr catalyzes the deubiquitination of bot specific gene promoters by activator nuclear receptor-mediated transactiv 58kDa and 60 kDa. The antibody is sp Author Pu Dianyun Ren 34 Zelong Han 36 Qingxiang Liu 29 Storage: Storage Buffer: PBS with 0.02% sodium azide and 50	ranscription regul th histones H2A a s such as MYC, wh ration and cell cy pecific to USP22. bmed ID 743406 333288 732405 ter shipment.	latory histone acety and H2B, thereby acti here it is required fo ccle progression. USF Journal Mol Oncol Cell Death Discov Sci Adv	lation (HAT) complex SAGA. USP22 ing as a coactivator. It is recruited to r transcription. USP22 is required for 22 has 2 isoforms with molecular mass Application WB,IP WB,IHC,IF,IP		
Notable Publications	human, mouse, rat USP22, also named as KIAA 1063 and deubiquitinating component of the tr catalyzes the deubiquitination of bot specific gene promoters by activator nuclear receptor-mediated transactiv 58kDa and 60 kDa. The antibody is sp Author Pu Dianyun Ren 34 Zelong Han 36 Qingxiang Liu 29 Storage: Storage Store at -20°C. Stable for one year aft Storage Buffer:	ranscription regul th histones H2A a s such as MYC, wh ration and cell cy pecific to USP22. bmed ID 743406 333288 732405 ter shipment.	latory histone acety and H2B, thereby acti here it is required fo ccle progression. USF Journal Mol Oncol Cell Death Discov Sci Adv	lation (HAT) complex SAGA. USP22 ing as a coactivator. It is recruited to r transcription. USP22 is required for 22 has 2 isoforms with molecular mass Application WB,IP WB,IHC,IF,IP		


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 55110-1-AP (USP22 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 55110-1-AP (USP22 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of USP22 antibody (55110-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-USP22 transfected A549 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 55110-1-AP (USP22 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



IP result of anti-USP22 (IP:55110-1-AP, 4ug; Detection:55110-1-AP 1:500) with HepG2 cells lysate 4000 ug.