#### For Research Use Only

# FOXC1 Polyclonal antibody

Catalog Number: 55365-1-AP

Featured Product

6 Publications



**Basic Information** 

Catalog Number: 55365-1-AP

Size:

GenBank Accession Number:

NM\_001453

GeneID (NCBI):

150ul, Concentration: 1000 ug/ml by 2296 Nanodrop and 740 ug/ml by Bradford UNIPROT ID:

method using BSA as the standard;

Source:

Rabbit Isotype: IgG

57 kDa Observed MW:

Q12948

Full Name:

forkhead box C1

Calculated MW:

70 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB 1:200-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

**Applications** 

**Tested Applications:** 

WB, IP, ELISA

Cited Applications: WB, IHC, CoIP, IF

Species Specificity: human, mouse

**Cited Species:** 

human, mouse

**Positive Controls:** 

WB: MCF-7 cells, HEK-293 cells, HepG2 cells

IP: HepG2 cells,

## **Background Information**

FOXC1, also named as FKHL7 and FREAC3, binding of FREAC-3 and FREAC-4 to their cognate sites results in bending of the DNA at an angle of 80-90 degrees. Defects in FOXC1 are the cause of Axenfeld-Rieger syndrome type 3 (RIEG3). Defects in FOXC1 are the cause of iridogoniodysgenesis anomaly (IGDA). Defects in FOXC1 are a cause of Peters anomaly. This antibody is specific to FOXC1. Phosphorylation modification of Foxc1 protein may be  $responsible \ for \ the \ larger \ molecular \ weight \ of \ detection \ compared \ with \ theoretical \ molecular \ weight$ (PMID:27708239;16403239).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Qinchen Cao	30431099	Mol Med Rep	WB
Yu-Zhong Xu	28635400	Tumour Biol	WB
Ya Wang	34461284	Bone	WB

Storage

Storage:

Store at -20°C.


Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

### Selected Validation Data







WB result of FOXC1 antibody (55365-1-AP; 1:300; incubated at room temperature for 1.5 hours) with sh-Control and sh-FOXC1 transfected HEK-293 cells.

IP result of anti-FOXC1 (IP:55365-1-AP, 4ug; Detection:55365-1-AP 1:300) with HepG2 cells lysate 4000 ug.

MCF-7 cells were subjected to SDS PAGE followed by western blot with 55365-1-AP (FOXC1 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.