

For Research Use Only

BMP6 Polyclonal antibody

Catalog Number: 55421-1-AP

3 Publications



Basic Information

Catalog Number: 55421-1-AP	GenBank Accession Number: NM_001718	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 800 ug/ml by Nanodrop;	GeneID (NCBI): 654	Recommended Dilutions: WB 1:500-1:1000 IHC 1:500-1:2000
Source: Rabbit	UNIPROT ID: P22004	
Isotype: IgG	Full Name: bone morphogenetic protein 6	
	Calculated MW: 57 kDa	
	Observed MW: 57 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls:
Cited Applications: WB, IHC	WB : A549 cells, mouse liver tissue, HEK-293 cells, HL-60 cells, U2OS cells
Species Specificity: human, mouse	IHC : human tonsillitis tissue, human liver tissue
Cited Species: human, mouse, rat	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Bone morphogenetic protein 6 (BMP6) is also named as VG-1-related protein (VG-1-R, VGR-1) and VGR, and belongs to the TGF-beta family (PMID:33878365). Growth factor of the TGF-beta superfamily plays essential roles in many developmental processes including cartilage and bone formation (PMID:31019025). BMP6 participates in iron homeostasis, inhibits invasion by increasing adhesions and cell-cell type interactions and induces angiogenesis directly on vascular endothelial cells. BMP6 is coded by a tumor suppressor gene whose subexpression is related to the development and cancer progression; during neoplastic processes, methylation is the main mechanism by which gene silencing occurs (PMID:34706618). BMP6 plays an important role in the regulation of hepcidin/HAMP expression and iron metabolism by acting as a ligand for hemojuvelin/HJV (PMID:26582087). BMP6 initiates the canonical BMP signaling cascade by associating with type I receptor ACVR1 and type II receptor ACVR2B (PMID:18070108).

Notable Publications

Author	Pubmed ID	Journal	Application
Yaoxue Guo	39797928	J Neuroimmune Pharmacol	WB
Kaiwei Wang	38485350	J Pharmacol Sci	IHC
Xuan Liu	38403744	Cell Tissue Res	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 55421-1-AP (BMP6 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 55421-1-AP (BMP6 antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 55421-1-AP (BMP6 antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse liver tissue were subjected to SDS PAGE followed by western blot with 55421-1-AP (BMP6 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.