

For Research Use Only

UBE2E1 Polyclonal antibody

Catalog Number: 55457-1-AP

Featured Product



Basic Information

Catalog Number: 55457-1-AP	GenBank Accession Number: NM_003341	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 550 µg/ml by Nanodrop;	GeneID (NCBI): 7324	Recommended Dilutions: WB 1:500-1:2400 IHC 1:50-1:500 IF 1:50-1:500
Source: Rabbit	Full Name: ubiquitin-conjugating enzyme E2E 1 (UBC4/5 homolog, yeast)	
Isotype: IgG	Calculated MW: 21 kDa	
	Observed MW: 21-27 kDa	

Applications

Tested Applications: IF, IHC, WB, ELISA	Positive Controls: WB : HEK-293 cells, IHC : human breast cancer tissue, IF : MCF-7 cells,
Species Specificity: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

UBE2E1 is a class III E2 ubiquitin conjugating enzyme that belongs to the UBE2E family of E2s, comprising UBE2E1/UBCH6, UBE2E2/UBCH8, and UBE2E3/UBCH9. UBE2E family members share a highly conserved UBC domain but are distinguished from one another by their unique N-terminal extensions. UBE2E1 contains a 50-residue unstructured N-terminal extension rich in serine and lysine residues, which has been shown to interact with USP7 and is essential for USP7-mediated UbE2E1 stabilization. In addition to its ubiquitin conjugating activity, UBE2E1 can also act as an ISG15 conjugating enzyme in vitro, although this activity has not been shown to occur in vivo. UBE2E1 can be covalently modified with either ISG15 or ubiquitin, both of which interfere with the ubiquitin conjugating activity of UBE2E1. UBE2E1 has also been reported to monoubiquitinate histone H2A at K119 in concert with the PRC1 E3 ligase complex. Full-length UBE2E1 is reported to be a dimer, with dimerization mediated by interaction of residues 21-27 from one molecule with the core domain of another molecule, as well as by contacts between a1 of the two interacting molecules. (PMID: 28073915, PMID: 30282802, PMID: 23871895)

Storage


Storage:
Store at -20°C.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of UBE2E1 antibody (55457-1-AP; 1:1200; incubated at room temperature for 1.5 hours) with sh-Control and sh-UBE2E1 transfected HEK-293 cells.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 55457-1-AP (UBE2E1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 55457-1-AP (UBE2E1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using 55457-1-AP (UBE2E1 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).