For Research Use Only

TGFBI / BIGH3 Monoclonal antibody

Catalog Number:60007-1-lg Featured Product 9 Publications




Basic Information	Catalog Number: GenBank Accession Number: 60007-1-lg BC000097		nber:	Purification Method: Protein A purification	
	Size:	GeneID (NCBI):		CloneNo.:	
	150ul, Concentration: 1772 ug/ml by	7045		3E11D11	
	Nanodrop and 1000 ug/ml by Bradfor method using BSA as the standard;	^d ENSEMBL Gene ID: ENSG00000120708		Recommended Dilutions: WB: 1:500-1:2000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC: 1:500-1:2000 IF-P: 1:200-1:800	
	Source: Mouse	UNIPROT ID: Q15582			
	Isotype: IgG2a Immunogen Catalog Number: AG0241	Full Name: transforming growth factor, beta- induced, 68kDa			
		Calculated MW: 683 aa, 75 kDa			
		Observed MW: 68 kDa			
Applications	Tested Applications:		Positive Controls:		
	WB, IHC, IF-P, IP, ELISA		VB : human kidney tissue,		
	Cited Applications: WB, IHC, IF, IP		IP : HeLa cells,		
	Species Specificity: IHC : human human			colon tissue, human skin cancer tissue, cancer tissue, human placenta tissue, v tissue	
	Cited Species:			colon cancer tissue,	
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen			
	TGFBI, also named as BIGH3, Kerato-epithelin and RGD-CAP, binds to type I, II, and IV collagens. TGFBI is an adhesion protein which may play an important role in cell-collagen interactions. In cartilage, it may be involved in endochondral bone formation. TGFBI is an extracellular matrix adaptor protein, it has been reported to be differentially expressed in transformed tissues. TGFBI is a predictive factor of the response to chemotherapy, and suggest the use of TGFBI-derived peptides as possible therapeutic adjuvants for the enhancement of responses to chemotherapy.(PMID:20509890) Defects in TGFBI are the cause of epithelial basement membrane corneal dystrophy (EBMD). Defects in TGFBI are the cause of corneal dystrophy Groenouw type 1 (CDGG1). Defects in TGFBI are the cause of corneal dystrophy lattice type 1 (CDL1). Defects in TGFBI are a cause of corneal dystrophy Thiel-Behnke type (CDTB). Defects in TGFBI are the cause of Reis-Buecklers corneal dystrophy (CDRB). Defects in TGFBI are the cause of lattice corneal dystrophy type 3A (CDL3A). Defects in TGFBI are the cause of Avellino corneal dystrophy (ACD).				
Background Information	endochondral bone formation. TGFBI differentially expressed in transform suggest the use of TGFBI-derived pep chemotherapy.(PMID:20509890) Defe (EBMD). Defects in TGFBI are the caus cause of corneal dystrophy lattice typ type (CDTB). Defects in TGFBI are the cause of lattice corneal dystrophy typ	is an extracellular matri ed tissues. TGFBI is a pre- tides as possible therape ects in TGFBI are the caus e of corneal dystrophy G e 1 (CDL1). Defects in TC cause of Reis-Buecklers	x adaptor pro dictive factor eutic adjuvan e of epithelia roenouw type ¡FBI are a cau corneal dystr	ctions. In cartilage, it may be involved i tein, it has been reported to be of the response to chemotherapy, and ts for the enhancement of responses to al basement membrane corneal dystrop 1 (CDGG1). Defects in TGFBI are the se of corneal dystrophy Thiel-Behnke ophy (CDRB). Defects in TGFBI are the	
	endochondral bone formation. TGFBI differentially expressed in transform suggest the use of TGFBI-derived pep chemotherapy.(PMID:20509890) Defe (EBMD). Defects in TGFBI are the caus cause of corneal dystrophy lattice typ type (CDTB). Defects in TGFBI are the cause of lattice corneal dystrophy typ (ACD).	is an extracellular matri ed tissues. TGFBI is a pre- tides as possible therape ects in TGFBI are the caus e of corneal dystrophy G e 1 (CDL1). Defects in TC cause of Reis-Buecklers	x adaptor pro dictive factor eutic adjuvan e of epithelia roenouw type iFBI are a cau corneal dystr TGFBI are th	ctions. In cartilage, it may be involved i tein, it has been reported to be of the response to chemotherapy, and ts for the enhancement of responses to al basement membrane corneal dystrop 1 (CDGG1). Defects in TGFBI are the se of corneal dystrophy Thiel-Behnke ophy (CDRB). Defects in TGFBI are the	
	endochondral bone formation. TGFBI differentially expressed in transform suggest the use of TGFBI-derived pep chemotherapy.(PMID:20509890) Defe (EBMD). Defects in TGFBI are the caus cause of corneal dystrophy lattice typ type (CDTB). Defects in TGFBI are the cause of lattice corneal dystrophy typ (ACD).	is an extracellular matri ed tissues. TGFBI is a pre- tides as possible therape ects in TGFBI are the caus e of corneal dystrophy G ie 1 (CDL1). Defects in TC cause of Reis-Buecklers ie 3A (CDL3A). Defects in	x adaptor pro dictive factor eutic adjuvan e of epithelia roenouw type iFBI are a cau corneal dystr TGFBI are th	ttions. In cartilage, it may be involved i tein, it has been reported to be of the response to chemotherapy, and ts for the enhancement of responses to al basement membrane corneal dystrop e 1 (CDGG1). Defects in TGFBI are the se of corneal dystrophy Thiel-Behnke ophy (CDRB). Defects in TGFBI are the e cause of Avellino corneal dystrophy	
	endochondral bone formation. TGFBI differentially expressed in transform suggest the use of TGFBI-derived pep chemotherapy.(PMID:20509890) Defe (EBMD). Defects in TGFBI are the caus cause of corneal dystrophy lattice typ type (CDTB). Defects in TGFBI are the cause of lattice corneal dystrophy typ (ACD). Author Put Laura S M Lecker 345	is an extracellular matri ed tissues. TGFBI is a pre- tides as possible therape eds in TGFBI are the caus e of corneal dystrophy G ee 1 (CDL1). Defects in TC cause of Reis-Buecklers ee 3A (CDL3A). Defects in med ID Journal	x adaptor pro dictive factor eutic adjuvan e of epithelia roenouw type iFBI are a cau corneal dystr .TGFBI are th	ttions. In cartilage, it may be involved it tein, it has been reported to be of the response to chemotherapy, and ts for the enhancement of responses to al basement membrane corneal dystrop et (CDGG1). Defects in TGFBI are the se of corneal dystrophy Thiel-Behnke ophy (CDRB). Defects in TGFBI are the e cause of Avellino corneal dystrophy Application	
	endochondral bone formation. TGFBI differentially expressed in transform suggest the use of TGFBI-derived pep chemotherapy.(PMID:20509890) Defe (EBMD). Defects in TGFBI are the caus cause of corneal dystrophy lattice typ type (CDTB). Defects in TGFBI are the cause of lattice corneal dystrophy typ (ACD). Author Put Laura S M Lecker 345 Tianhong Pan 291	is an extracellular matri ed tissues. TGFBI is a pre- tides as possible therape cts in TGFBI are the caus e of corneal dystrophy G ee 1 (CDL1). Defects in TC cause of Reis-Buecklers is 3A (CDL3A). Defects in med ID Journal 661272 Cancer	x adaptor pro dictive factor eutic adjuvan e of epithelia roenouw type iFBI are a cau corneal dystr TGFBI are th L Res	ttions. In cartilage, it may be involved it tein, it has been reported to be of the response to chemotherapy, and ts for the enhancement of responses to al basement membrane corneal dystrop e 1 (CDGG1). Defects in TGFBI are the se of corneal dystrophy Thiel-Behnke ophy (CDRB). Defects in TGFBI are the e cause of Avellino corneal dystrophy Application WB,IHC	
Background Information	endochondral bone formation. TGFBI differentially expressed in transform suggest the use of TGFBI-derived pep chemotherapy.(PMID:20509890) Defe (EBMD). Defects in TGFBI are the caus cause of corneal dystrophy lattice typ type (CDTB). Defects in TGFBI are the cause of lattice corneal dystrophy typ (ACD). Author Put Laura S M Lecker 345 Tianhong Pan 291 Douglas S Annis 262 Storage: Storage Storage Store at -20°C. Stable for one year aft Storage Buffer:	is an extracellular matri ed tissues. TGFBI is a pre- tides as possible therape cts in TGFBI are the caus e of corneal dystrophy G use 1 (CDL1). Defects in TC cause of Reis-Buecklers is 3A (CDL3A). Defects in med ID Journal i61272 Cancer .90493 Neopla 173833 PLoS O	x adaptor pro dictive factor eutic adjuvan e of epithelia roenouw type iFBI are a cau corneal dystr TGFBI are th L Res	ttions. In cartilage, it may be involved i tein, it has been reported to be of the response to chemotherapy, and ts for the enhancement of responses to al basement membrane corneal dystrop e 1 (CDGG1). Defects in TGFBI are the se of corneal dystrophy Thiel-Behnke pophy (CDRB). Defects in TGFBI are the e cause of Avellino corneal dystrophy Application WB,IHC WB,IHC	
Notable Publications	endochondral bone formation. TGFBI differentially expressed in transform suggest the use of TGFBI-derived pep chemotherapy.(PMID:20509890) Defe (EBMD). Defects in TGFBI are the caus cause of corneal dystrophy lattice typ type (CDTB). Defects in TGFBI are the cause of lattice corneal dystrophy typ (ACD). Author Put Laura S M Lecker 345 Tianhong Pan 291 Douglas S Annis 262 Storage: Store at -20°C. Stable for one year aft	is an extracellular matri ed tissues. TGFBI is a pre- tides as possible therape cts in TGFBI are the caus e of corneal dystrophy G use 1 (CDL1). Defects in TC cause of Reis-Buecklers is 3A (CDL3A). Defects in med ID Journal i61272 Cancer .90493 Neopla .73833 PLoS O er shipment.	x adaptor pro dictive factor eutic adjuvan e of epithelia roenouw type iFBI are a cau corneal dystr TGFBI are th L Res	ttions. In cartilage, it may be involved tein, it has been reported to be of the response to chemotherapy, and ts for the enhancement of responses to al basement membrane corneal dystrop e 1 (CDGG1). Defects in TGFBI are the se of corneal dystrophy Thiel-Behnke ophy (CDRB). Defects in TGFBI are the e cause of Avellino corneal dystrophy Application WB,IHC WB,IHC	

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human kidney tissue were subjected to SDS PAGE followed by western blot with 60007-1-1g (TGFBI / BIGH3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 60007-1-Ig (TGFBI / BIGH3 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 60007-1-Ig (TGFBI / BIGH3 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-TGFBI / BIGH3 (IP:60007-1-Ig, 4ug; Detection:60007-1-Ig 1:300) with HeLa cells lysate 1200ug.



Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using TGFBI / BIGH3 antibody (60007-1-Ig, Clone: 3E11D11) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using TGFBI / BIGH3 antibody (60007-1-Ig, Clone: 3E11D11) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).