

For Research Use Only

CXCR3B-specific Monoclonal antibody, PBS Only



Catalog Number: 60065-1-PBS

Featured Product

Basic Information

Catalog Number:

60065-1-PBS

Size:

100ug, Concentration: 1 mg/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG2b

GenBank Accession Number:

AF469635

GeneID (NCBI):

2833

UNIPROT ID:

P49682

Full Name:

chemokine (C-X-C motif) receptor 3

Calculated MW:

46 kDa

Observed MW:

43 kDa

Purification Method:

Protein A purification

CloneNo.:

1B2D6

Applications

Tested Applications:

WB, FC, IHC, Indirect ELISA

Species Specificity:

human

Background Information

CXCR3 (C-X-C chemokine receptor type 3) is a G protein-coupled, seven-transmembrane domain chemokine receptor that is expressed on the surface of a number of cell types, including activated CD4+ and CD8+ T cells, NK and NK-T cells, plasmacytoid dendritic cells, and some B cells (PMID: 16847335). CXCR3 binds and is activated by the three IFN-gamma-inducible chemokines: CXCL9, CXCL10, and CXCL11. Binding of these ligands to CXCR3 has been implicated in regulating T-cell infiltration during inflammation and tissue injury (PMID: 17250586). Alternatively spliced transcript variants encoding different isoforms have been found for this gene. One of the isoforms, CXCR3B, mediates the angiostatic activity of CXCR3 ligands and also acts as functional receptor for CXCL4 (PMID:12782716). This monoclonal antibody is specific to CXCR3B.

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




1X10⁶ U-937 cells were intracellularly stained with 0.4 ug Anti-Human CXCR3B-specific (60065-1-Ig, Clone:1B2D6) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 60065-1-PBS in a



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60065-1-Ig (CXCR3B-specific antibody at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60065-1-PBS in a different storage buffer formulation.



MOLT-4 cells were subjected to SDS PAGE followed by western blot with 60065-1-Ig (CXCR3B-specific antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60065-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60065-1-Ig (CXCR3B-specific antibody at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60065-1-PBS in a different storage buffer formulation.