For Research Use Only

RRM1 Monoclonal antibody

Catalog Number:60073-2-lg Featured Product

6 Publications



Basic Information

Catalog Number: GenBank Accession Number:

60073-2-lg BC006498 GeneID (NCBI): Size: 150ul, Concentration: 2000 ug/ml by 6240

Nanodrop and 1000 ug/ml by Bradford_{UNIPROT ID:} method using BSA as the standard; P23921

Source: Full Name:

Mouse ribonucleotide reductase M1 Isotype: Calculated MW:

90 kDa Immunogen Catalog Number: Observed MW:

AG0789 90 kDa

Tested Applications:

lgG2b

WB, IHC, IF/ICC, IF-P, IP, ELISA

Cited Applications: WB, IHC, IF

Species Specificity:

human **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Purification Method: Protein A purification

CloneNo.: 5H6F3

Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:3000-1:8000 IF-P 1:200-1:800 IF/ICC 1:400-1:1600

Applications

Positive Controls:

WB: K-562 cells, HeLa cells

IP: K-562 cells.

IHC: human breast cancer tissue, human colon cancer tissue, human lung cancer tissue, human pancreas cancer tissue, human urothelial carcinoma tissue

IF-P: human breast cancer tissue. IF/ICC: HepG2 cells, HeLa cells

Background Information

Ribonucleoside-diphosphate reductase functions as a heterodimer of a large and a small subunits in deoxyribonucleotide synthesis. RRM1 constitutes to the large subunit (R1) of ribonucleotide reductase, and it can either form heterodimer with small subunit RRM or RRM2B(PMID:16376858). RRM1 provides the precursors necessary for DNA synthesis. RRM1 can not be detected in quiescent cells, while its mRNA and protein are present throughout the cell cycle in cycling cells(PMID:8188248). Researches showed that RRM1 is involved in carcinogenesis, tumor progression, and the resistance of non-small-cell lung cancer (NSCLC) to treatment. Low level expression of RRM1 in NSCLC is associated with poor survival (PMID:17314339).

Notable Publications

Author	Pubmed ID	Journal	Application
Toru Aoyama	28521448	Oncol Lett	IHC
Donghua Geng	35837166	J Gastrointest Oncol	WB
Sean G Rudd	31950591	EMBO Mol Med	WB

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




K-562 cells were subjected to SDS PAGE followed by western blot with 60073-2-lg (RRM1 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60073-2-lg (RRM1 antibody) at dilution of 1:5000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60073-2-lg (RRM1 antibody) at dilution of 1:5000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using RRM1 antibody (60073-2-1g, Clone: 5H6F3) at dilution of 1:400 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using RRM1 antibody (60073-2-lg, Clone: 5H6F3) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



IP result of anti-RRM1 (IP:60073-2-Ig, 5ug; Detection:60073-2-Ig 1:1000) with K-562 cells lysate 3440ug.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 60073-2-Ig (RRM1 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).