## For Research Use Only

## RRM1 Monoclonal antibody, PBS Only

Catalog Number:60073-2-PBS Featured Product



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

60073-2-PBS

GeneID (NCBI):

Protein A purification

BC006498

CloneNo.: 5H6F3

100ug, Concentration: 1mg/ml by

**UNIPROT ID:** 

Nanodrop:

P23921 Full Name:

Mouse Isotype:

ribonucleotide reductase M1

lgG2b

Calculated MW: 90 kDa

Immunogen Catalog Number: AG0789

Observed MW:

90 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IF-P, IP, Indirect ELISA

Species Specificity:

human

## **Background Information**

Ribonucleoside-diphosphate reductase functions as a heterodimer of a large and a small subunits in deoxyribonucleotide synthesis. RRM1 constitutes to the large subunit (R1) of ribonucleotide reductase, and it can either form heterodimer with small subunit RRM or RRM2B(PMID:16376858). RRM1 provides the precursors necessary for DNA synthesis. RRM1 can not be detected in quiescent cells, while its mRNA and protein are present throughout the cell cycle in cycling cells(PMID:8188248). Researches showed that RRM1 is involved in carcinogenesis, tumor progression, and the resistance of non-small-cell lung cancer (NSCLC) to treatment. Low level expression of RRM1 in NSCLC is associated with poor survival (PMID:17314339).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


## Selected Validation Data




K-562 cells were subjected to SDS PAGE followed by western blot with 60073-2-lg (RRM1 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60073-2-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60073-2-lg (RRM1 antibody) at dilution of 1:5000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60073-2-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60073-2-lg (RRM1 antibody) at dilution of 1:5000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60073-2-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using RRM1 antibody (60073-2-lg, Clone: 5H6F3) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60073-2-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using RRM1 antibody (60073-2-Ig, Clone: 5H6F3) at dilution of 1:800 and Coralite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60073-2-PBS in a different storage buffer formulation.



IP result of anti-RRM1 (IP:60073-2-Ig, 5ug; Detection:60073-2-Ig 1:1000) with K-562 cells lysate 3440ug. This data was developed using the same antibody clone with 60073-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 60073-2-lg (RRM1 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60073-2-PBS in a different storage buffer formulation