## For Research Use Only

## IRF5 Monoclonal antibody

Catalog Number:60088-1-lg 1 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** Protein A purification

60088-1-lg

GeneID (NCBI):

150ul , Concentration: 500  $\mu g/ml$  by

CloneNo.: 2A10C5

Bradford method using BSA as the

**UNIPROT ID:** 

BC004201

Recommended Dilutions:

standard;

Q13568

WB 1:500-1:2000

Source:

Full Name:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Mouse Isotype:

interferon regulatory factor 5 Calculated MW:

protein lysate

IgG2a

56 kDa

IHC 1:50-1:500 IF/ICC 1:20-1:200

Immunogen Catalog Number:

Observed MW:

AG0841

63+110 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IP, ELISA

Positive Controls: WB: HeLa cells, HepG2 cells

Cited Applications: WB

IP: HeLa cells.

IF/ICC: HepG2 cells,

Species Specificity:

IHC: human placenta tissue, human breast cancer

human, mouse, pig

tissue, human tonsillitis tissue

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Background Information** 

IRF5, also named as SLEB10, contians one IRF tryptophan pentad repeat DNA-binding domain and belongs to the IRF family. It is a transcription factor involved in the induction of interferons IFNA and INFB and inflammatory cytokines upon virus infection. It is activated by TLR7 or TLR8 signaling. Genetic variations in IRF5 are associated with susceptibility to inflammatory bowel disease type 14 (IBD14) and systemic lupus erythematosus type 10  $^{\circ}$ (SLEB10).

**Notable Publications** 

Author	Pubmed ID	Journal	Application
Martin Heather J HJ	17609264	J Virol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




HeLa cells were subjected to SDS PAGE followed by western blot with 60088-1-1g (IRF5 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 burs.




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 60088-1-Ig (IRF5 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 60088-1-1g (IRF5 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-IRF5 (IP:60088-1-Ig. 4ug; Detection:60088-1-Ig 1:500) with HeLa cells lysate 1600ug.



Immunofluorescent analysis of HepG2 cells using 60088-1-Ig (IRF5 antibody) at dilution of 1:50 and Rhodamine-Goat anti-Mouse IgG.