For Research Use Only

Human IgA Monoclonal antibody

Catalog Number:60099-1-lg 4 Publications



Basic Information

Catalog Number: GenBank Accession Number:

60099-1-lg BC016369
Size: GeneI D (NCBI):
150ul , Concentration: 1900 ug/ml by 3493

Nanodrop and 1000 ug/ml by Bradford_{Full Name}:

method using BSA as the standard; immunoglobulin heavy constant

Source: alpha 1
Mouse Calculated MW:

Isotype:496 aa, 53 kDaIgG1Observed MW:Immunogen Catalog Number:55-60 kDa

AG7417

Purification Method: Protein A purification

CloneNo.: 3A5A5

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:160000-1:640000 IF-P 1:200-1:800

Applications

Tested Applications: WB, IHC, IF-P, ELISA

Cited Applications:

IF

Species Specificity:

human Cited Species: rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: human saliva tissue, human saliva, human

plasma

IHC: human tonsillitis tissue,
IF-P: human tonsillitis tissue,

Background Information

Ig alpha is the major immunoglobulin class in body secretions. It may serve both to defend against local infection and to prevent access of foreign antigens to the general immunologic system. This antibody recognizes the Chain C region of human Ig alpha-1 (IGHA1).

Notable Publications

Author	Pubmed ID	Journal	Application
Zhipeng Li	30670042	J Nanobiotechnology	
Felix Marsh-Wakefield	39761010	Hepatol Commun	
Wu Gu	39248343	Adv Sci (Weinh)	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human saliva were subjected to SDS PAGE followed by western blot with 60099-1-Ig (Human IgA Antibody) at dilution of 1:16000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using Human IgA antibody (60099-1-Ig, Clone: 3A5A5) at dilution of 1:400 and CoraLite®488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60099-1-lg (Human IgA antibody) at dilution of 1:320000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60099-1-Ig (Human IgA antibody) at dilution of 1:50000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).