## For Research Use Only

## CXCL8/IL-8 Monoclonal antibody

Catalog Number:60141-2-lg 6 Publications



**Basic Information** 

Catalog Number: 60141-2-lg

GenBank Accession Number:

BC013615

GeneID (NCBI):

Full Name:

interleukin 8 Calculated MW:

99 aa. 11 kDa

150ul, Concentration: 1000 µg/ml by 3576

Bradford method using BSA as the

standard:

Mouse Isotype:

IgG2a

Immunogen Catalog Number:

AG10552

**Purification Method:** Protein A purification

CloneNo.: 3D11D5

**Applications** 

**Tested Applications:** 

**Cited Applications:** ELISA, IHC, WB

Species Specificity:

human

human

**Cited Species:** 

## **Background Information**

Interleukin 8 (IL-8), also known as CXCL8, which is a member of the CXC chemokine family. This chemokine is secreted by a variety of cell types including monocyte/macrophages, T cells, neutrophils, fibroblasts, endothelial cells, and various tumor cell lines in response to inflammatory stimuli. IL-8 has two primary functions. It induces chemotaxis in target cells, primarily neutrophils but also other granulocytes, causing them to migrate toward the site of infection. IL-8 also induces phagocytosis once they have arrived. This gene is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. IL-8 is also known to be a potent promoter of angiogenesis. IL-8 has been associated with tumor angiogenesis, metastasis, and poor prognosis in breast cancer. IL-8 may present a novel therapeutic target for estrogen driven breast carcinogenesis and tumor progression. The human IL-8 cDNA sequence predicts a protein of 99 amino acids. Removal of a 22-residue signal peptide generates a mature protein of 77 amino acids (~8 kDa).

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Qixia Xu	31493351	Aging Cell	WB
Changxu Wang	36202915	Oncogene	WB
Sen Wang	28745433	Cancer Med	IHC

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

Selected Validation Data