

FGF12 Monoclonal antibody

Catalog Number: 60152-1-Ig

Basic Information

Catalog Number: 60152-1-Ig	GenBank Accession Number: BC022524	Purification Method: Caprylic Acid/Ammonium Sulfate Precipitation
Size: 150ul , Concentration: 1200 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2257	CloneNo.: 2F11A10
Source: Mouse	Full Name: fibroblast growth factor 12	Recommended Dilutions: WB 1:500-1:2000
Isotype: IgG2b	Calculated MW: 181 aa, 20 kDa	
Immunogen Catalog Number: AG5046	Observed MW: 20 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : Human brain, tissue
Species Specificity: human	

Background Information

FGF12 is a member of the fibroblast growth factor (FGF) family. FGF family members play important roles in embryogenesis, angiogenesis, and wound repair. FGF12 lacks the N-terminal signal sequence present in most of the FGF family members, but it contains clusters of basic residues that have been demonstrated to act as a nuclear localization signal. When transfected into mammalian cells, this protein accumulated in the nucleus, but was not secreted. FGF12 plays an intracellular role in the inhibition of radiation-induced apoptosis. FGF12 is expressed abundantly in developing and adult nervous systems; therefore, FGF12 was thought to be related to nervous system development and function.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 60152-1-Ig (FGF12 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.