For Research Use Only

TEM1 Monoclonal antibody

Catalog Number:60170-1-lg 7 Publications




Basic Information	Catalog Number: 60170-1-lg	GenBank Accession Number BC051340	er: Purification Method: Protein A purification
	Size:	GenelD (NCBI):	CloneNo.:
	150ul , Concentration: 1600 ug/ml by		1F9B4
	Nanodrop and 1000 ug/ml by Bradford		Recommended Dilutions:
	method using BSA as the standard;	Q9HCU0	WB 1:1000-1:4000
	Source:	Full Name:	IHC 1:50-1:500
	Mouse CD248 molecule, endosialin		
	lsotype: lgG1	Calculated MW: 757 aa, 81 kDa	
	Immunogen Catalog Number: AG13334	Observed MW: 150-160 kDa	
Applications	Tested Applications:	Pos	itive Controls:
	WB, IHC, ELISA	WB	: HUVEC cells, COLO 320 cells, bEnd.3 cells,
	Cited Applications:	NIF	I/3T3 cells
	WB, IHC, IF, IP, CoIP	IHC	: human breast cancer tissue, human renal cell
	Species Specificity: human, mouse	car	cinoma tissue
	Cited Species: human, mouse, rat		
	Note-IHC: suggested antigen ro TE buffer pH 9.0; (*) Alternativ retrieval may be performed wa buffer pH 6.0	ely, antigen	
Background Information	(CTLD) containing type I transmembra activated perivascular and stromal ce plays a role in tumor angiogenesis. It malignant disease. Two isoforms of h	ane glycoprotein. It is now o lls, detected in most cancer is a potential diagnostic to uman TEM1 exist. The calcu	sialin and CD164L1, is a C-type lectin-like domain considered to be a highly selective marker for 's and at least some inflammatory disorders. CD24 ol and therapeutic target of inflammatory and ulated molecular weights of the two isoforms are 8 and the glycosylated form has a larger apparent
	(CTLD) containing type I transmembra activated perivascular and stromal ce plays a role in tumor angiogenesis. It malignant disease. Two isoforms of h kDa and 46 kDa, respectively. Native molecular weight than 81 kDa.	ane glycoprotein. It is now o Ils, detected in most cancer is a potential diagnostic to uman TEM1 exist. The calco TEM1 can be glycosylated,	considered to be a highly selective marker for rs and at least some inflammatory disorders. CD24 ol and therapeutic target of inflammatory and llated molecular weights of the two isoforms are 8 and the glycosylated form has a larger apparent
	(CTLD) containing type I transmembra activated perivascular and stromal ce plays a role in tumor angiogenesis. It malignant disease. Two isoforms of h kDa and 46 kDa, respectively. Native molecular weight than 81 kDa.	ane glycoprotein. It is now o lls, detected in most cancer is a potential diagnostic to uman TEM1 exist. The calco TEM1 can be glycosylated, med ID Journal	considered to be a highly selective marker for rs and at least some inflammatory disorders. CD24 ol and therapeutic target of inflammatory and ulated molecular weights of the two isoforms are 8 and the glycosylated form has a larger apparent Application
	(CTLD) containing type I transmembra activated perivascular and stromal ce plays a role in tumor angiogenesis. It malignant disease. Two isoforms of h kDa and 46 kDa, respectively. Native molecular weight than 81 kDa. Author Pub Shengya Cao 345	ane glycoprotein. It is now of Ils, detected in most cancer is a potential diagnostic to uman TEM1 exist. The calco TEM1 can be glycosylated, med ID Journal 31301 Proc Natl /	considered to be a highly selective marker for rs and at least some inflammatory disorders. CD24 ol and therapeutic target of inflammatory and ulated molecular weights of the two isoforms are 8 and the glycosylated form has a larger apparent Acad Sci U S A WB
Background Information	(CTLD) containing type I transmembra activated perivascular and stromal ce plays a role in tumor angiogenesis. It malignant disease. Two isoforms of h kDa and 46 kDa, respectively. Native molecular weight than 81 kDa.AuthorPut Shengya CaoShengya Cao345Chia-Lun Hong359	ane glycoprotein. It is now of lls, detected in most cancer is a potential diagnostic to uman TEM1 exist. The calcu TEM1 can be glycosylated, med ID Journal 31301 Proc Natl / 50912 Cancer Re	Acad Sci U S A Solution S
	(CTLD) containing type I transmembra activated perivascular and stromal ce plays a role in tumor angiogenesis. It malignant disease. Two isoforms of h kDa and 46 kDa, respectively. Native molecular weight than 81 kDa.AuthorPut Shengya CaoShengya Cao345Chia-Lun Hong359	ane glycoprotein. It is now of Ils, detected in most cancer is a potential diagnostic to uman TEM1 exist. The calco TEM1 can be glycosylated, med ID Journal 31301 Proc Natl /	considered to be a highly selective marker for rs and at least some inflammatory disorders. CD24 ol and therapeutic target of inflammatory and ulated molecular weights of the two isoforms are 8 and the glycosylated form has a larger apparent Acad Sci U S A WB
Notable Publications	(CTLD) containing type I transmembra activated perivascular and stromal ceplays a role in tumor angiogenesis. It malignant disease. Two isoforms of h kDa and 46 kDa, respectively. Native molecular weight than 81 kDa. Author Put Shengya Cao 345 Chia-Lun Hong 359 Po-Sheng Chen 357 Storage: Storage Buffer: PBS with 0.02% sodium azide and 50%	ane glycoprotein. It is now of lls, detected in most cancer is a potential diagnostic to uman TEM1 exist. The calco TEM1 can be glycosylated, med ID Journal 31301 Proc Natl A 150912 Cancer Re 32643 Sci Rep er shipment. % glycerol pH 7.3.	Acad Sci U S A Solution S
	(CTLD) containing type I transmembra activated perivascular and stromal ceplays a role in tumor angiogenesis. It malignant disease. Two isoforms of h kDa and 46 kDa, respectively. Native molecular weight than 81 kDa. Author Pub Shengya Cao 345 Chia-Lun Hong 359 Po-Sheng Chen 357 Storage: Storage Storage Buffer: Storage Buffer:	ane glycoprotein. It is now of lls, detected in most cancer is a potential diagnostic to uman TEM1 exist. The calco TEM1 can be glycosylated, med ID Journal 31301 Proc Natl A 150912 Cancer Re 32643 Sci Rep er shipment. % glycerol pH 7.3.	Acad Sci U S A Solution Sci U S A WB,IF,IP Solution Sci U S A Solution Sci U S A

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





HUVEC cells were subjected to SDS PAGE followed by western blot with 60170-1-1g (TEM1 antibody at dilution of 1:2000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60170-1-1g (TEM1 antibody at dilution of 1:250 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60170-1-1g (TEM1 antibody at dilution of 1:250 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).