

For Research Use Only

CD122/IL2RB Monoclonal antibody

Catalog Number: 60174-1-Ig



Basic Information

Catalog Number: 60174-1-Ig	GenBank Accession Number: BC025691	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 3560	CloneNo.: 3A12C7
Source: Mouse	Full Name: interleukin 2 receptor, beta	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Calculated MW: 551 aa, 61 kDa	
Immunogen Catalog Number: AG3929	Observed MW: 75 kDa, 61 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : NK-92 cells,
Species Specificity: human	

Background Information

The interleukin 2 receptor, which is involved in T cell-mediated immune responses, is present in 3 forms with respect to ability to bind interleukin 2. The low affinity form is a monomer of the alpha subunit and is not involved in signal transduction. The intermediate affinity form consists of an alpha/beta subunit heterodimer, while the high affinity form consists of an alpha/beta/gamma subunit heterotrimer. The beta subunit (IL2RB, also known as CD122 and p75) is involved in receptor mediated endocytosis and transduces the mitogenic signals of nterleukin 2.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.1% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Untreated and PNGase F-treated lysates of NK-92 cells were subjected to SDS PAGE followed by western blot with 60174-1-Ig (CD122/IL2RB antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.