For Research Use Only

CD34 Monoclonal antibody, PBS Only

Catalog Number:60180-2-PBS



Purification Method:

CloneNo.:

4F11H3

Protein G purification

Basic Information

Catalog Number: 60180-2-PBS

Nanodrop;

GenBank Accession Number:

BC039146

GeneID (NCBI):

Size: 100ug, Concentration: 1mg/ml by

UNIPROT ID:

P28906 Source Mouse Full Name: Isotype: CD34 molecule lgG1 Calculated MW:

Immunogen Catalog Number: 41 kDa

AG5996 Observed MW:

105 kDa

Applications

Tested Applications:

WB, IHC, Indirect ELISA

Species Specificity:

human


Background Information

CD34 is a 105- to 120-kDa glycophosphoprotein expressed on the majority of hematopoietic stem/progenitor cells, bone marrow stromal cells, capillary endothelial cells, embryonic fibroblasts, and some nerve tissue. CD34 is a $commonly\ used\ marker\ for\ identifying\ human\ hematopoietic\ stem/progenitor\ cells\ and\ mediates\ cell\ adhesion\ adhesion\$ lymphocyte homing by binding L-selectin and E-selectin ligands. CD34 is also one of the best negative selection markers for characterizing and/or isolating human MSCs from bone marrow and other sources. Along with other positive selection markers (such as CD29, CD44, CD90, CD105 and CD166), negative selection markers (such as CD34 and CD45) are used for MSC identification. The calculated molecular mass of human CD34 is 41 kDa, various forms with different molecular weights may be produced due to different glycosylation patterns and alternative splicing (PMID: 24375067; 15750786).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data



human placenta tissue were subjected to SDS PAGE followed by western blot with 60180-2-lg (CD34 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60180-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60180-2-lg (CD34 antibody) at dilution of 1:10000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60180-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60180-2-lg (CD34 antibody) at dilution of 1:10000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60180-2-PBS in a different storage buffer formulation.

Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60180-2-lg (CD34 antibody) at dilution of 1:20000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60180-2-PBS in a different storage buffer formulation.