For Research Use Only

NF-L Monoclonal antibody, PBS Only

Catalog Number: 60189-1-PBS



Purification Method:

CloneNo.:

5C12G4

Protein A purification

Basic Information

Catalog Number: GenBank Accession Number:

60189-1-PBS BC039237

GeneID (NCBI):

100ug, Concentration: 1mg/ml by

Nanodrop: **UNIPROT ID:** P07196 Mouse Full Name:

Isotype: neurofilament, light polypeptide

lgG1 Calculated MW: Immunogen Catalog Number: 543 aa, 62 kDa AG15178 Observed MW:

65 kDa

Applications

Tested Applications:

WB, IHC, IF-P, FC (Intra), Indirect ELISA

Species Specificity: human, mouse, rat, pig


Background Information

 $NEFL, also \ named \ as \ NF68 \ and \ NF-L, belongs \ to \ the \ intermediate \ filament \ family. \ Neurofilaments \ are \ the \ 10 \ nm$ intermediate filaments found specifically in neurons. They are a major component of the cell's cytoskeleton, and provide support for normal axonal radial growth. Neurofilaments usually contain three intermediate filament proteins: L, M, and H, which are involved in the maintenance of neuronal caliber. The names given to the three major neurofilament subunits are based upon the apparent molecular weight of the mammalian subunits on SDS-PAGE: NF-L, 65-68 kDa; NF-M,145-160 kDa and NF-H, 200-220 kDa. This antibody is specific to NEFL.


Storage

Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 60189-1-lg (NF-L antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60189-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 60189-1-lg (NF-L antibody) at dilution of 1:100000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60189-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using 60189-1-1g (NF-L antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60189-1-PBS in a different storage buffer formulation.



1X10^6 PC-12 cells were intracellularly stained with 0.4 ug Anti-Human NF-L (60189-1-1g, Clone:5C12G4) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 60189-1-PBS in a different storage buffer formulation.

Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 60189-1-lg (NF-L antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60189-1-PBS in a different storage buffer formulation.