For Research Use Only

# TNFR1 Monoclonal antibody, PBS Only



Catalog Number:60192-1-PBS

**Featured Product** 

### **Basic Information**

Catalog Number: 60192-1-PBS Size: 100ug , Concentration: 1mg/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG16112 GenBank Accession Number: BC010140 GeneID (NCBI): 7132 UNIPROT ID: P19438 Full Name: tumor necrosis factor receptor superfamily, member 1A Calculated MW: 455 aa, 50 kDa Observed MW:

50-55 kDa

Purification Method: Protein A purification CloneNo.: 2A6E3

Applications

Tested Applications: WB, IF, IHC, ELISA Species Specificity: human, mouse, rat, pig

#### **Background Information**

Tumor necrosis factor (TNF) is a multifunctional cytokine that plays a key role in regulating inflammation, immune functions, host defense, and apoptosis (PMID: 16407280). TNF exists in soluble and membrane-bound forms. TNF signals through two distinct cell surface receptors, TNFR1 (TNFRSF1A, CD120a) and TNFR2 (TNFRSF1B, CD120b). Whereas TNFR1 is widely expressed, expression of TNFR2 is limited to cells of the immune system, endothelial cells, and nerve cells (PMID: 22053109). TNFR1, which contains a death domain (DD) within its intracytoplasmic region, is thought to be the key receptor for TNF signaling (PMID: 16407280). This receptor can activate NF-kappaB, mediate apoptosis, and function as a regulator of inflammation. Antiapoptotic protein BCL2-associated athanogene 4 (BAG4/SODD) and adaptor proteins TRADD and TRAF2 have been shown to interact with this receptor, and thus play regulatory roles in the signal transduction mediated by the receptor.


#### Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60192-1-Ig (TNFR1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 60192-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using TNFR1 antibody (60192-1-lg, Clone: 2A6E3) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60192-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 60192-1-1g (TNFR1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60192-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 60192-1-Ig (TNFR1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60192-1-PBS in a different storage buffer formulation.