

For Research Use Only

AKT Monoclonal antibody

Catalog Number: 60203-2-Ig

Featured Product

547 Publications



Basic Information

Catalog Number:

60203-2-Ig

Size:

150ul, Concentration: 800 ug/ml by Bradford method using BSA as the standard;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG16695

GenBank Accession Number:

BC000479

GeneID (NCBI):

207

UNIPROT ID:

P31749

Full Name:

v-akt murine thymoma viral oncogene homolog 1

Calculated MW:

56 kDa

Observed MW:

56-62 kDa

Purification Method:

Protein A purification

CloneNo.:

2C5D1

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:100-1:400

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, rabbit, chicken, bovine, hamster, goat, sheep, duck

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, rat liver tissue, mouse brain tissue, HEK-293 cells, NIH/3T3 cells, RAW 264.7 cells, ROS1728 cells, LNCaP cells, Hela cells, Jurkat cells, HSC-T6 cells, PC-12 cells

IP: mouse brain tissue,

IHC: human breast cancer tissue, human cervical cancer tissue

IF/ICC: MCF-7 cells,

Background Information

The serine-threonine protein kinase AKT1 is catalytically inactive in serum-starved primary and immortalized fibroblasts. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. This antibody detects all the members of AKT with/without phospho- modification.

Notable Publications

Author	Pubmed ID	Journal	Application
Yi Yu	34585393	J Periodontol	WB
Wenbin Pei	34650433	Front Pharmacol	WB
YanHua Fan	36174847	Fitoterapia	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60203-2-Ig (AKT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




WB result of AKT antibody (60203-2-Ig; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-AKT transfected A549 cells.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 60203-2-Ig (AKT Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using AKT antibody (60203-2-Ig, Clone: 2C5D1) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 60203-2-Ig (AKT Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HEK-293 cells were subjected to SDS PAGE followed by western blot with 60203-2-Ig (AKT Antibody) and 66444-1-Ig (AKT1-phospho-S473 Antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



IP result of anti-AKT (IP:60203-2-Ig, 5ug; Detection:60203-2-Ig 1:1000) with mouse brain tissue lysate 4000ug.