For Research Use Only

Galectin-3 Monoclonal antibody

Catalog Number:60207-1-lg Featured Product

32 Publications



Basic Information

Catalog Number: GenBank Accession Number: **Purification Method:** 60207-1-lg BC001120 Protein A purification

GeneID (NCBI): CloneNo.: 150ul, Concentration: 1000 µg/ml by 3958 1C1B2

Recommended Dilutions: lectin, galactoside-binding, soluble, 3 WB 1:5000-1:50000 Source: IHC 1:1000-1:4000 Mouse Calculated MW: IF 1:400-1:1600 Isotype: 26 kDa

lgG2b Observed MW: Immunogen Catalog Number: 31 kDa

AG6753

Applications

Tested Applications:

FC, IF, IHC, WB, ELISA Cited Applications: IF, IHC, IP, WB Species Specificity: human, mouse, rat **Cited Species:**

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, HSC-T6 cells, HeLa cells, A549 cells, MCF-7 cells, T-47D cells, HepG2 cells, Caco-2 cells, human placenta tissue

IHC: human thyroid cancer tissue, human colon tissue,

human ovary tumor tissue

IF: HeLa cells, mouse kidney tissue, MCF-7 cells, human thyroid cancer tissue

Background Information

Galectins are a family of animal lectins defined by shared characteristic amino-acid sequences and affinity for β galactose-containing oligosac-charides (PMID: 8063692). Galectin-3, a member of the β -galactoside-binding proteins, contains one carbohydrate recognition domain (CRD) and a proline- and glycine-rich N-terminal domain through which is able to form oligomers (PMID: 14758078). Galectin-3 is widely expressed in many normal tissues and a variety of tumors. It is found intracellularly in nucleus and cytoplasm or secreted outside of cell, being present on the cell surface or in the extracellular space (PMID: 16478649). Galectin-3 is involved in various biological processes including cell growth, adhesion, differentiation, apoptosis, angiogenesis, immune response, neoplastic transformation and metastasis (PMID: 16478649; 14758078).

Notable Publications

Author	Pubmed ID	Journal	Application
Vinay V Eapen	34585663	Elife	WB,IF
Dandan Zong	36332381	Sleep Med	WB
Ying Li	33124742	FASEBJ	WB

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60207-1-lg (Galectin-3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




WB result of Galectin-3 specific antibody (60207-1-lg, 1:1000) with si-control and si-Galectin-3 transfected HeLa cells.




Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 60207-1-Ig (Galectin-3 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 60207-1-Ig (Galectin-3 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using Galectin-3 antibody (60207-1-Ig. Clone: 1C1B2) at dilution of 1:800 and CoraLite®594-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).



1X10^6 HeLa cells were intracellularly stained with 0.8 ug Anti-Human Galectin-3 (60207-1-lg, Clone:1C1B2) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.8 ug Mouse IgG2b isotype control Mouse McAb (66360-3-lg, Clone: 11B8C4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).