For Research Use Only

# CD7 Monoclonal antibody, PBS Only

Catalog Number:60209-2-PBS



### **Basic Information**

- Catalog Number: 60209-2-PBS Size: 100ug , Concentration: 1mg/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG1805
- GenBank Accession Number: BC009293 GeneID (NCBI): 924 ENSEMBL Gene ID: ENSG00000173762 UNIPROT ID: P09564 Full Name: CD7 molecule Calculated MW: 240 aa, 25 kDa Observed MW: 34-36 kDa

#### Purification Method: Protein G purification CloneNo.: 2A4E6

## **Applications**

Tested Applications: WB, IHC, Indirect ELISA Species Specificity: human, mouse

### **Background Information**

CD7 is a type I transmembrane glycoprotein expressed on thymocytes and mature T cells. It is the most reliable clinical marker of T-cell acute lymphocytic leukemia.


### Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

### Selected Validation Data




Untreated and PNGase F-treated lysates of Jurkat cells were subjected to SDS PAGE followed by western blot with 60209-2-Ig (CD7 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. PNGase F was obtained from Atagenix (cat.NO. ata808). This data was developed using the same antibody clone with 60209-2-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 60209-2-1g (CD7 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60209-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60209-2-1g (CD7 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60209-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60209-2-1g (CD7 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60209-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60209-2-Ig (CD7 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60209-2-PBS in a different storage buffer formulation.