For Research Use Only

BCS1L Monoclonal antibody

Catalog Number:60212-1-lg Featured Product



Basic Information

Applications

Catalog Number: GenBank Accession Number:

60212-1-lg BC007500 GeneID (NCBI): Size:

150ul, Concentration: 1100 ug/ml by 617 Nanodrop and 860 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; Q9Y276

Source: Full Name: Mouse BCS1-like (yeast) Isotype: Calculated MW: lgG1 48 kDa Immunogen Catalog Number: Observed MW:

AG18280 47 kDa

Tested Applications:

WB, IHC, IF/ICC, ELISA Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Protein G purification CloneNo.:

3G6H1

Positive Controls:

human kidney tissue

IF/ICC: MCF-7 cells,

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:20-1:200 IF/ICC 1:20-1:200

WB: Colo320 cells, COLO 320 cells, HEK-293 cells

IHC: human gliomas tissue, human brain tissue,

Background Information

Human BCS1-like (BCS1L), a mitochondrial inner-membrane protein, is a chaperone necessary for the assembly of mitochondrial respiratory chain complex III. Recently studies indicated that the mutations in this protein may cause the GRACILE (growth retardation, aminoaciduria, cholestasis, iron overload, lactacidosis, and early death) syndrome, a recessively inherited lethal disease characterized by fetal growth retardation, lactic acidosis, aminoaciduria, cholestasis, and abnormalities in iron metabolism.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




WB result of BCS1L antibody (60212-1-lg; 1:300; incubated at room temperature for 1.5 hours) with sh-Control and sh-BCS1L transfected HEK-293 cells.




COLO 320 cells were subjected to SDS PAGE followed by western blot with 60212-1-1g (BCS1L antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas using 60212-1-Ig(BCS1L antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human gliomas using 60212-1-Ig(BCS1L antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of () fixed MCF-7 cells using 60212-1-Ig (BCS1L antibody) at dilution of 1:50 and Rhodamine-Goat anti-Rabbit IgG.