## For Research Use Only

# CD9 Monoclonal antibody

Catalog Number:60232-1-lg

**Featured Product** 

76 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

60232-1-lg BC011988 Protein G purification
Size: GenelD (NCBI): CloneNo.:

150ul , Concentration: 1500 µg/ml by 928

Nanodrop;
Full Name:

Source:
CD9 molecule
Mouse
Calculated MW:
Isotype:
228 aa, 25 kDa
IgG1
Observed MW:
Immunogen Catalog Number:
23-27 kDa

AG14529

**Applications** 

**Tested Applications:** 

FC, IF, IHC, WB, ELISA Cited Applications: FC, IF, IHC, PLA, WB Species Specificity: human

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, HeLa cells

IHC: human ovary tumor tissue, human colon cancer tissue, human tonsillitis tissue, human breast cancer tissue

**Purification Method:** 

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:1000-1:4000

IF 1:200-1:800

4H7B9

IF: human breast cancer tissue, human ovary tumor tissue, human lung cancer tissue

# **Background Information**

The cell-surface molecule CD9, a member of the transmembrane-4 superfamily, interacts with the integrin family and other membrane proteins, and is postulated to participate in cell migration and adhesion. Expression of CD9 enhances membrane fusion between muscle cells and promotes viral infection in some cells (PMID:10459022). It is often used as a mesenchymal stem cell marker (PMID:18005405). CD9 is also known as the p24 antigen besides MIC3, TSPAN29 because it is a protein of molecular weight 24 kD. The CD9 antigen appears to be a 227-amino acid molecule with 4 hydrophobic domains and 1 N-glycosylation site.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Kosuke Otani	31561474	Int J Mol Sci	WB
Na-Na Sun	34483252	Chin Med J (Engl)	WB
Zhi-Hong Zong	31666098	J Exp Clin Cancer Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## Selected Validation Data




WB result of CD9 antibody (60232-1-Ig, 1:2000) with si-Control and si-CD9 transfected HeLa cells.




Various lysates were subjected to SDS PAGE followed by western blot with 60232-1-lg (CD9 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 60232-1-lg (CD9 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 60232-1-lg (CD9 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using CD9 antibody (60232-1-Ig, Clone: 4H7B9) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using CD9 antibody (60232-1-Ig, Clone: 4H7B9) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10^6 HeLa cells were stained with 0.2 ug Anti-Human CD9 (60232-1-1g, Clone:4H7B9) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (green), or 0.2 ug isotype control (black). Cells were fixed with 4%