For Research Use Only

IL-17F Monoclonal antibody, PBS Only



Purification Method:

Protein A purification

CloneNo.:

6H5F10

Catalog Number: 60245-1-PBS

Basic Information

Catalog Number:

GenBank Accession Number:

60245-1-PBS

Nanodrop:

Mouse

Isotype:

AG16310

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by

BC070124

112744

UNIPROT ID:

Q96PD4

Full Name: interleukin 17F

lgG1 Calculated MW: Immunogen Catalog Number: 163 aa, 18 kDa

Observed MW:

37 kDa

Applications

Tested Applications:

WB, Indirect ELISA

Species Specificity:

human, mouse

Background Information

The interleukin 17 (IL-17) family of cytokines contains 6 structurally related cytokines, IL-17A, IL-17B, IL-17C, IL-17D, IL-17E and IL-17F. IL-17 family plays crucial roles in host defense against microbial organisms and in the development of inflammatory diseases. IL-17A is a pro-inflammatory cytokine that also has the capacity to promote $angiogenesis \ and \ osteoclastogenesis. \ IL-17F \ shares \ the \ highest \ homology \ with \ IL-17A \ and \ signals \ via \ a \ receptor$ composed by the IL-17RA and IL-17RC subunits. IL-17A and IL-17F can form IL-17A/A or IL-17F/F homodimers, IL-17A/F heterodimers are also formed. IL-17A and IL-17F, produced by the Th17 CD4(+) T cell lineage, have been linked to a variety of inflammatory and autoimmune conditions. IL-17F levels are elevated in sera and lesional psoriatic skin compared to non-lesional tissue. IL-17F also has been implicated in the development of neutrophilic airway inflammation.

Storage

Storage:

Store at -80°C. Storage Buffer:

PBS Only

Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 60245-1-lg (IL-17F antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60245-1-PBS in a different storage buffer formulation.