## For Research Use Only

## CD20 Monoclonal antibody, PBS Only

Catalog Number: 60271-1-PBS



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** Protein A purification

60271-1-PBS

GeneID (NCBI):

CloneNo.:

100ug, Concentration: 1mg/ml by

BC002807

4A7G3

Nanodrop;

**UNIPROT ID:** P11836

Mouse Isotype:

Full Name: membrane-spanning 4-domains,

lgG2b

subfamily A, member 1

Immunogen Catalog Number: AG17309

Calculated MW: 297 aa, 33 kDa

Observed MW: 35 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IF-P, Indirect ELISA

Species Specificity:

**Background Information** 


CD20 is a 33-37 kDa transmembrane phosphoprotein. CD20 is a B-lymphocyte surface molecule that is widely expressed during B-cell ontogeny, from early pre-B-cell developmental stages until final differentiation into plasma  $cells. \ CD20 \ functions \ as \ calcium-permeable \ cation \ channel. \ It \ is \ involved \ in \ the \ regulation \ of \ B-cell \ activation \ and \ cells \ activation \ and \ cells \ ce$ proliferation. CD20 serves as a useful target for antibody-mediated therapeutic depletion of B-cells.

Storage


Storage:

Store at -80°C. Storage Buffer: PBS Only


## Selected Validation Data




Raji cells (CD20+), Daudi cells (CD20+), Ramos cells (CD20+), Jurkat cells (CD20-), and HeLa cells (CD20-) were subjected to SDS PAGE followed by western blot with 60271-1-1g (CD20 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60271-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60271-1-lg (CD20 antibody) at dilution of 1:12800 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60271-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60271-1-lg (CD20 antibody) at dilution of 1:12800 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60271-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD20 antibody (60271-1-lg, Clone: 4A7G3) at dilution of 1:400 and CoraLite® 594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CD3 antibody (17617-1-AP, green). This data was developed using the same antibody clone with 60271-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsilliris tissue using CD20 antibody (60271-1-1g, Clone: 4A7G3) at dilution of 1:400 and CoraLite® 594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CD3 antibody (17617-1-AP, green). This data was developed using the same antibody clone with 60271-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed Raji cells using CD20 antibody (60271-1-1g, Clone: 4A7G3) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 60271-1-PBS in a different storage buffer formulation.

Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60271-1-lg (CD20 antibody) at dilution of 1:20000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60271-1-PBS in a different storage buffer formulation.