For Research Use Only

PLAP Monoclonal antibody, PBS Only

Catalog Number:60294-1-PBS Featured Product



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

60294-1-PBS

BC009647 GeneID (NCBI): Protein G purification

CloneNo.: 2B7C9

100ug, Concentration: 1mg/ml by Nanodrop;

UNIPROT ID:

Mouse

P05187

Isotype:

Full Name: alkaline phosphatase, placental

lgG1

(Regan isozyme)

Immunogen Catalog Number:

Calculated MW:

AG13361

58 kDa

Observed MW: 66 kDa

Applications

Tested Applications:

WB, IHC, Indirect ELISA

Species Specificity:

Background Information

Placental alkaline phosphatase (PLAP) is a glycosyl phosphatidylinositol (GPI)-anchored sialoglycoprotein, which $shows\ greater\ resistance\ to\ heat\ inactivation\ than\ other\ alkaline\ phosphatase.\ Presented\ at\ high\ levels\ in\ placental\ placenta$

trophoblasts, PLAP is recognized as an important marker of differentiation in human malignancies.

Storage


Storage:

Store at -80°C.


Storage Buffer:

PBS Only


Selected Validation Data




WB result of PLAP antibody (60294-1-Ig, 1:1000) with si-control and si-PLAP transfected HepG2 cells. This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.




HepG2 cells were subjected to SDS PAGE followed by western blot with 60294-1-1g (PLAP antibody at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 60294-1-1g (PLAP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.




human placenta tissue were subjected to SDS PAGE followed by western blot with 60294-1-lg (PLAP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 60294-1-lg (PLAP antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 60294-1-lg (PLAP antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 60294-1-lg (PLAP antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation



Immunohistochemical analysis of paraffinembedded human malignant melanoma tissue slide using 60294-1-lg (PLAP antibody) at dilution of 1:4000 (under 10x lens). This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human malignant melanoma tissue slide using 60294-1-lg (PLAP antibody) at dilution of 1:4000 (under 40x lens). This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.