

For Research Use Only

HSP90 Monoclonal antibody

Catalog Number: 60318-1-Ig **93 Publications**



Basic Information

Catalog Number: 60318-1-Ig	GenBank Accession Number: BC023006	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 3320	CloneNo.: 3F11C1
Source: Mouse	UNIPROT ID: P07900	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG2a	Full Name: heat shock protein 90kDa alpha (cytosolic), class A member 1	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
Immunogen Catalog Number: AG3826	Calculated MW: 853 aa, 90 kDa	IHC 1:50-1:500
	Observed MW: 85-90 kDa	IF/ICC 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, IP, CoIP, ELISA	Positive Controls: WB : LNCaP cells, HeLa cells, HEK-293 cells, MCF-7 cells, Jurkat cells, HSC-T6 cells, ROS1728 cells, NIH/3T3 cells, 4T1 cells
Cited Applications: WB, IHC, IF, IP, CoIP, RIP	IP : HepG2 cells,
Species Specificity: human, mouse, rat, pig	IHC : human ovary tumor tissue,
Cited Species: human, mouse, rat, pig, monkey, chicken	IF/ICC : HepG2 cells,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

HSP90, encoded by HSP90AA1, is a constitutively and ubiquitously expressed molecular chaperone that is crucial for the stability and function of many proteins. HSP90 provides chaperoning activity for client proteins; many of them are members of oncogenic pathways, indicating its implication in tumor malignancy. HSP90 mainly resides in the cytosol, while it can also be released to the extracellular space. Secreted Hsp90 is a C-terminal truncated form. It has been reported that the level of plasma Hsp90 is positively correlated with tumor malignancy in clinical cancer patients, and can be a promising diagnostic marker for tumor malignancy in clinical application.

Notable Publications

Author	Pubmed ID	Journal	Application
Yunfei Chen	32966240	Aging (Albany NY)	WB
Xingfeng Liu	36131205	Nat Metab	WB
Rui Lin	36082450	Prostate	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Co-IP result of anti-HSP90 (IP:60318-1-Ig, 4ug; Detection:13422-1-AP 1:5000) with mouse brain tissue lysate 2240ug.




Various lysates were subjected to SDS PAGE followed by western blot with 60318-1-Ig (HSP90 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.




Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 60318-1-Ig (HSP90 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 60318-1-Ig (HSP90 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 60318-1-Ig (HSP90 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



IP result of anti-HSP90 (IP:60318-1-Ig, 5ug; Detection:60318-1-Ig 1:5000) with HepG2 cells lysate 1200 ug.