For Research Use Only

P-selectin / CD62P Monoclonal antibody



Catalog Number:60322-1-lg

20 Publications

Basic Information

Catalog Number: GenBank Accession Number: **Purification Method:** 60322-1-lg BC028067 Protein A purification

Size: GeneID (NCBI): CloneNo.: 150ul, Concentration: 1295 µg/ml by 6403 3G2C7

Nanodrop and 1080 µg/ml by Bradford_{Full Name}: Recommended Dilutions: method using BSA as the standard;

selectin P (granule membrane protein WB 1:2000-1:10000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

140kDa, antigen CD62) Mouse protein lysate Calculated MW:

830 aa, 91 kDa Isotype: lgG2b Observed MW: Immunogen Catalog Number: 140 kDa, 105-110 kDa

AG4124

Applications Tested Applications:

IP. WB. FLISA

Cited Applications: FC, IF, IHC, WB Species Specificity:

human **Cited Species:**

human, rat, mouse

Positive Controls:

WB: human peripheral blood platelets, Jurkat cells

IP: K-562 cells,

Background Information

P-selectin, also known as GMP140 or CD62P, is a transmembrane glycoprotein that mediates the interaction of activated endothelial cells or platelets with leukocytes. It is an adhesion molecule involved in the pathogenesis of inflammation, thrombosis, and oncogenesis. P-selectin is stored in the alpha-granules of platelets and Weibel-Palade bodies of endothelial cells. Upon cell activation by agonists, P-selectin is transported rapidly to the cell surface.

Notable Publications

Author	Pubmed ID	Journal	Application
Jinwei Pang	34566870	Front Neurol	WB,IF
LinXi Cheng	34489520	Commun Biol	FC
Chunyu Huang	33015023	Front Bioeng Biotechnol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Untreated and PNGase F-treated lysates of human platelets were subjected to SDS PAGE followed by western blot with 60322-1-1g (P-selectin / CD62P antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. PNGase F was obtained from Atagenix (cat.NO. ata808).



human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 60322-1-lg (P-selectin / CD62P antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP Result of anti-P-selectin / CD62P (IP:60322-1-Ig, 5ug; Detection:60322-1-Ig 1:600) with K-562 cells lysate 1760ug.