## For Research Use Only

## Perilipin-2 Monoclonal antibody

Mouse

Catalog Number: 60340-1-lg



**Basic Information** 

Catalog Number: GenBank Accession Number:

60340-1-lg BC005127 GeneID (NCBI):

150ul, Concentration: 1000 ug/ml by 123 Nanodrop: **UNIPROT ID:** Q99541

Isotype: adipose differentiation-related

lgG1 protein

Immunogen Catalog Number: Calculated MW: AG7643 48 kDa

> Observed MW: 47-52 kDa

Full Name:

**Applications** 

**Tested Applications:** WB, IHC, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human adipose tissue.

IHC: human renal cell carcinoma tissue, mouse brown adipose tissue, rat brown adipose tissue

**Purification Method:** 

Protein A purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:200-1:8000

CloneNo.:

1G5D4

## **Background Information**

ADRP (adipocyte differentiation related protein) also known as ADFP, adipophilin, or perilipin-2, is a member of PAT family which is responsible for the transportation of lipids and the formation of lipid droplets. ADRP is localized on the surface of lipid droplets in a variety of tissues and cell lines. ADRP is not detected in undifferentiated cells but increases rapidly to high levels when adipocyte precursors differentiate into adipocytes. Anti-ADRP antibody is a reliable and sensitive marker for lipid droplet. Enhanced expression of ADRP is linked to diseases with abnormal lipid storage, including hepatic steatosis, atherosclerosis and diabetes. Immunohistochemistry of ADRP may facilitate histomorphological diagnosis of these diseases.

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**



human adipose tissue were subjected to SDS PAGE followed by western blot with 60340-1-lg (ADRP Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brown adipose slide using 60340-1-Ig (ADRP/Perilipin 2 antibody) at dilution of 1:16000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-1g (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-1g (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).