

For Research Use Only

TrkA Monoclonal antibody, PBS Only

Catalog Number: 60548-1-PBS



Basic Information

Catalog Number: 60548-1-PBS	GenBank Accession Number: NM_001007792	Purification Method: Protein G purification
Size: 100ug , Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 4914	CloneNo.: 3B11C7
Source: Mouse	UNIPROT ID: P04629	
Isotype: IgG1	Full Name: neurotrophic tyrosine kinase, receptor, type 1	
Immunogen Catalog Number: AG27985	Calculated MW: 88 kDa	
	Observed MW: 140 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human

Background Information

NTRK1 (Neurotrophic tyrosine kinase receptor 1) is located on chromosome 1q23.1 and is responsible for a rare genetic disorder, CIPA (congenital insensitivity to pain with anhidrosis) (PMID: 37907480). As a neurotrophin receptor, NTRK1 is abundantly expressed in human neuronal tissues and plays an integral role in the development and function of the human nervous system (PMID: 27843590). NTRK1 also regulates the survival and proliferation of cholinergic neurons (PMID: 20054812).

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 60548-1-Ig (NTRK1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60548-1-PBS in a different storage buffer formulation.



NK-92 cells were subjected to SDS PAGE followed by western blot with 60548-1-Ig (NTRK1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60548-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 60548-1-Ig (NTRK1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60548-1-PBS in a different storage buffer formulation.