

For Research Use Only

HSD17B4 Monoclonal antibody

Catalog Number:60951-1-Ig



Basic Information

| | | |
|--|---|--|
| Catalog Number: 60951-1-Ig | GenBank Accession Number: BC003098 | Purification Method: Protein A purification |
| Size: 150ul , Concentration: 1000 ug/ml by Nanodrop; | GeneID (NCBI): 3295 | CloneNo.: 1A3H11 |
| Source: Mouse | UNIPROT ID: P51659 | Recommended Dilutions: WB: 1:5000-1:50000 IF/ICC: 1:1000-1:4000 |
| Isotype: IgG2a | Full Name: hydroxysteroid (17-beta) dehydrogenase 4 | |
| Immunogen Catalog Number: AG7221 | Calculated MW: 80 kDa | |
| | Observed MW: 80 kDa, 45 kDa | |

Applications

| | |
|--|---|
| Tested Applications: WB, IF/ICC, ELISA | Positive Controls: |
| Species Specificity: human | WB : HeLa cells, LNCaP cells, HepG2 cells, RT-4 cells IF/ICC : HeLa cells, |

Background Information

HSD17B4 (17-beta-hydroxysteroid dehydrogenase 4) is also named as Peroxisomal multifunctional enzyme type 2, D-bifunctional protein or multifunctional protein 2. It codes for a 80 kDa enzyme containing three distinct functional domains and is localized in peroxisomes. It is a bifunctional enzyme acting on the peroxisomal beta-oxidation pathway for fatty acids and catalyzing the formation of 3-ketoacyl-CoA intermediates from both straight-chain and 2-methyl-branched-chain fatty acids. After peroxisomal import, the full-length protein is proteolytically cleaved to yield a 35-kDa dehydrogenase subunit and a 45-kDa hydratase subunit containing the hydratase and SCP domains (PMID: 28868548, 24602372).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 60951-1-Ig (HSD17B4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HSD17B4 antibody (60951-1-Ig, Clone: 1A3H11) at dilution of 1:2000 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-phalloidin (red).