#### For Research Use Only

# Anti-Mouse CD107a / LAMP1 (1D4B)

Catalog Number:65050-1-lg 4 Publications



**Basic Information** 

Catalog Number: 65050-1-lg

Size:

100ug , 0.5 mg/ml

Source:

Isotype:

IgG2a, kappa

16783

P11438

**UNIPROT ID:** 

Full Name:

protein 1

GenBank Accession Number: **Purification Method:** BC006785 Protein G purification GeneID (NCBI):

CloneNo.: 1D4B

Recommended Dilutions: IF/ICC 1:50-1:500

lysosomal-associated membrane

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra)

**Cited Applications:** 

Species Specificity:

mouse Cited Species:

mouse

**Positive Controls:** 

IF/ICC: NIH/3T3 cells,

### **Background Information**

LAMP1 (CD107a) is a heavily glycosylated membrane protein enriched in the lysosomal membrane. LAMP1 is extensively glycosylated with asparagine-linked oligosaccharides which protect it from intracellular proteolysis (PMID: 10521503). Although LAMP1 is expressed largely in the endosome-lysosomal membrane of cells, it is also found on the plasma membrane (PMID: 16168398). Elevated LAMP1 expression at the cell surface has also been detected during platelet and granulocytic cell activation, as well as in some tumor cells (PMID: 29085473). LAMP1 functions to provide selectins with carbohydrate ligands. This protein has also been shown to be a marker of degranulation on lymphocytes such as CD8+ and NK cells and may also play a role in tumor cell differentiation and metastasis (PMID: 18835598; 29085473; 9426697).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Panpan Wang	39183331	Mol Neurodegener	IF
Li Wang	38513294	J Leukoc Biol	IF
Weidong Le	37292937	Res Sq	IF

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

## **Selected Validation Data**



1X10^6 NIH/3T3 cells were intracellularly stained with 0.5 ug Anti-Mouse CD107a / LAMP1 (65050-1-Ig, Clone:1D4B) and FITC anti-Rat IgG2a antibody at dilution 1:100, or 0.5 ug Rat IgG2a Isotype Control (2A3) (65209-1-Ig, Clone: 2A3) and FITC anti-Rat IgG2a antibody at dilution 1:100. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using 65050-1-Ig (CD107a antibody) at dilution of 1:100 and Rhodamine (TRITC)-conjugated Goat Anti-Rat IgG(H+L).