For Research Use Only

Anti-Mouse CD31 (390)

Catalog Number:65058-1-lg 8 Publications



Purification Method:

Affinity purification

IF/ICC 1:50-1:500

Recommended Dilutions:

CloneNo.:

Basic Information

Catalog Number: 65058-1-lg

Size:

100ug , 0.5 mg/ml

Source:

Isotype:

IgG2a, kappa

18613

Q08481

platelet/endothelial cell adhesion

molecule 1

Applications

Tested Applications:

IF/ICC, FC

Cited Applications:

Species Specificity:

Mouse

Cited Species:

mouse

GenBank Accession Number:

BC008519 GeneID (NCBI):

UNIPROT ID:

Full Name:

Positive Controls:

IF/ICC: bEnd.3 cells,

Background Information

Platelet endothelial cell adhesion molecule-1 (PECAM-1, CD31) is a member of the immunoglobulin gene superfamily of cell adhesion molecules. CD31 is a transmembrane glycoprotein that is highly expressed on the surface of the endothelium, making up a large portion of its intracellular junctions. PECAM-1 is also present on the surface of hematopoietic cells and immune cells including platelets, monocytes, neutrophils, natural killer cells, megakaryocytes and some types of T-cell (PMID: 9011572). As well as its role in cell-cell adhesion, PECAM-1 functions as a signaling receptor, and is involved in important physiological events such as nitric oxide production, $regulation\ of\ T-cell\ immunity\ and\ tolerance, leukocyte\ transendothelial\ migration\ and\ inflammation\ and$ angiogenesis (PMID: 21183735; 20978210; 17872453; 20634489).

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaopeng Ai	35697191	J Ethnopharmacol	IF
Ziyi Kang	38838924	J Ethnopharmacol	IF
Jiaojiao Chen	38565041	Int Immunopharmacol	IF

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed bEnd.3 cells using CD31 antibody (65058-1-lg, Clone: 390) at dilution of 1:100 and CoraLite® 488-Conjugated Goat Anti-Rat IgG(H+L).



1x10^6 mouse splenocytes were surface stained with 0.5 ug Anti-Mouse CD31 (65058-1-lg, Clone: 390) and FITC anti-rat IgG2a Antibody at dilution 1:100 (red), or 0.5 ug Rat IgG2a Isotype Control (2A3) (65209-1-lg, Clone: 2A3) and FITC anti-rat IgG2a Antibody at dilution 1:100 (blue). Cells were not fixed.