For Research Use Only

Anti-Mouse CD8a (53-6.7) Catalog Number:65069-1-lg 5 Publications




Basic Information	Catalog Number: 65069-1-1g			Purification Method: Affinity purification
	Size: 500ug, 0.5 mg/ml	GeneID (NCB 12525	I):	CloneNo.: 53-6.7
	Source:	UNIPROT ID:		
	Rat	P01731		
	<mark>Isotype:</mark> IgG2a, kappa	Full Name: CD8 antigen,	alpha chain	
Applications	Tested Applications: FC			
	Cited Applications: FC			
	Species Specificity: mouse			
	Cited Species: mouse			
Background Information	CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8a β heterodimer while CD8a α homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) $\gamma \delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).			
0	been found on some gut int 2111591; 1831127; 842097 activation (PMID: 8499079)	raepithelial T cells , on s 5). CD8 acts as a co-rece	some T cell receptor (ptor that binds to MH	TCR) γδT cells and on NK cells (PMID: C class-I and participates in cytotoxicT ce
	been found on some gut int 2111591; 1831127; 842097 activation (PMID: 8499079) (PMID: 1968084).	raepithelial T cells , on s 5). CD8 acts as a co-rece . During T cell developm	some T cell receptor (ptor that binds to MH nent, CD8 is required	TCR) γδ T cells and on NK cells (PMID: C class-I and participates in cytotoxic T ce for positive selection of CD4-/CD8+ T cell
	been found on some gut int 2111591; 1831127; 842097. activation (PMID: 8499079) (PMID: 1968084). Author	raepithelial T cells , on s 5). CD8 acts as a co-rece . During T cell developm Pubmed ID	some T cell receptor (ptor that binds to MH nent, CD8 is required Journal	TCR) γδ T cells and on NK cells (PMID: C class-I and participates in cytotoxic T ce for positive selection of CD4-/CD8+ T cell Application
	been found on some gut int 2111591; 1831127; 842097 activation (PMID: 8499079) (PMID: 1968084). Author Huajing Wang	raepithelial T cells , on s 5). CD8 acts as a co-rece . During T cell developm Pubmed ID 34090420	some T cell receptor (ptor that binds to MH nent, CD8 is required Journal Malar J	TCR) γδ T cells and on NK cells (PMID: C class-I and participates in cytotoxic T ce for positive selection of CD4-/CD8+ T cell Application FC
Notable Publications	been found on some gut int 2111591; 1831127; 842097. activation (PMID: 8499079) (PMID: 1968084). Author	raepithelial T cells , on s 5). CD8 acts as a co-rece . During T cell developm Pubmed ID	some T cell receptor (ptor that binds to MH nent, CD8 is required Journal	TCR) γδ T cells and on NK cells (PMID: C class-I and participates in cytotoxic T ce for positive selection of CD4-/CD8+ T cell Application FC FC

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^{^6}C 57BL/6 mouse splenocytes were surface stained with CoraLite® Plus 647 Anti-Mouse CD3 (17A2) and 0.5 ug Anti-Mouse CD8a (65069-1-lg, Clone:53-6.7) or 0.5 ug Rat IgG2a Isotype Control (2A3) and FITC anti-rat IgG2a Antibody at dilution 1:100. Cells were not fixed.