

For Research Use Only

# Anti-Human ICAM-1/CD54 (15.2)

Catalog Number: 65075-1-Ig



## Basic Information

<b>Catalog Number:</b> 65075-1-Ig	<b>GenBank Accession Number:</b> BC015969	<b>Purification Method:</b> Affinity purification
<b>Size:</b> 100ug, 0.5 mg/ml	<b>GeneID (NCBI):</b> 3383	<b>CloneNo.:</b> 15.2
<b>Source:</b> Mouse	<b>ENSEMBL Gene ID:</b> ENSG00000090339	
<b>Isotype:</b> IgG1, kappa	<b>UNIPROT ID:</b> P05362	
	<b>Full Name:</b> intercellular adhesion molecule 1	
	<b>Calculated MW:</b> 90 kDa	

## Applications

**Tested Applications:**  
FC

**Species Specificity:**  
Human

## Background Information

ICAM-1 (CD54) is a 90-kDa transmembrane glycoprotein of the immunoglobulin superfamily and is critical for the firm attachment and transmigration of leukocytes out of blood vessels and into tissues (PMID: 19307690). ICAM-1 is expressed by several cell types, typically on endothelial cells and cells of the immune system, and its expression can be up-regulated by various stimuli, including TNF- $\alpha$ , INF- $\gamma$ , IL-1 and thrombin (PMID: 3086451; 9694714; 15979056). It is a ligand for LFA-1 and Mac-1, serves as a receptor for rhinovirus, and is one of several receptors used by Plasmodium falciparum (PMID: 2566624; 2538244; 2475784).

## Storage

**Storage:**  
Store at 2-8°C. Stable for one year after shipment.

**Storage Buffer:**  
PBS with 0.09% sodium azide.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10<sup>6</sup> human PBMCs were surface stained with 0.2 ug Anti-Human CD54 (ICAM-1) (65075-1-Ig, Clone:15.2) and APC-Donkey anti-Mouse IgG at dilution 1:1000. Cells were not fixed. Monocytes were gated.



1X10<sup>6</sup> human PBMCs were surface stained with 0.2 ug Anti-Human CD54 (ICAM-1) (65075-1-Ig, Clone:15.2) and APC-Donkey anti-Mouse IgG at dilution 1:1000. Cells were not fixed. Monocytes were gated.