For Research Use Only

## Anti-Mouse TCR beta (H57-597)

Catalog Number:65106-1-Ig



**Basic Information** 

Catalog Number: 65106-1-lg

....

100ug , 0.5 mg/ml

Source:

Armenian Hamster

Isotype: IgG GenBank Accession Number:

GeneID (NCBI):

21577 Full Name:

T-cell receptor beta chain

Purification Method: Affinity purification

CloneNo.: H57-597

**Applications** 

Tested Applications:

FC

Species Specificity:

Mouse

## **Background Information**


T-cell receptor (TCR) consists of heterodimeric glycoproteins (TCR alpha/beta or gamma/delta) that demonstrate homology with immunoglobulins (PMID: 2144901). TCR beta is a member of the immunoglobulin superfamily and is expressed on alpha/beta TCR-bearing T cells and thymocytes. The TCR heterodimers are noncovalently associated with CD3. The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta (PMID: 1826255). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Storage

Storage: Store at 2-8°C. Storage Buffer:

Phosphate based buffer with 0.09% sodium azide, pH 7.2.

## Selected Validation Data



1X10^6 mouse splenocytes were surface stained with 0.5 ug Anti-Mouse TCR beta (65106-1-1g, Clone:H57-597) and FITC Goat anti-hamster (Armenian) IgG Antibody at dilution 1:200 (green), or stained with 0.5 ug isotype control antibody and FITC Goat anti-hamster (Armenian) IgG Antibody at dilution 1:200 (black). Cells were not fixed.