For Research Use Only

Anti-Human CD11b (ICRF44)

Catalog Number:65116-1-lg 2 Publications



Basic Information

Catalog Number:

65116-1-lg

Size:

500ug , 0.5 mg/ml

Source: Mouse Isotype:

IgG1, kappa

ENSEMBL Gene ID: ENSG00000169896

GenBank Accession Number:

UNIPROT ID: P11215 Full Name:

BC096346

3684

GeneID (NCBI):

integrin, alpha M (complement component 3 receptor 3 subunit)

Calculated MW: 1152 aa, 127 kDa Purification Method: Affinity purification CloneNo.:

ICRF44

Applications

Tested Applications:

FC

Cited Applications:

FC

Species Specificity:

Human
Cited Species:
human

Background Information

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits (PMID: 9779984). CD11b, also known as Integrin alpha M or CR3A, belongs to the integrin alpha chain family. CD11b forms an α/β heterodimer with CD18 (integrin β 2). CD11b/CD18 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles and pathogens (PMID: 9558116; 20008295). CD11b/CD18 is a receptor for the complement protein fragment iC3b, and is also a receptor for fibrinogen, factor X and ICAM1 (PMID: 2971974; 15485828).

Notable Publications

| Author | Pubmed ID | Journal | Application |
|---------|-----------|---------------------|-------------|
| Hao Lei | 36173787 | J Med Chem | FC |
| Bin Qi | 33363140 | Front Cell Dev Biol | FC |

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1X10^6 human PBMCs were surface stained with 0.5 ug Anti-Human CD11b (65116-1-lg, Clone:ICRF 44) and Coralite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red) or unstained. Cells were not fixed. Monocytes were gated.



1X10^6 human PBMCs were surface stained with 0.5 ug Anti-Human CD11b (65116-1-lg, Clone:ICRF 44) and Coralite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 or unstained. Cells were not fixed. Monocytes were gated.