

For Research Use Only

Anti-Mouse PD-1/CD279 (J43.1)

Catalog Number: 65120-1-Ig



Basic Information

Catalog Number:	GenBank Accession Number:	Purification Method:
65120-1-Ig	BC119179	Affinity purification
Size:	GeneID (NCBI):	CloneNo.:
500ug , 0.5 mg/ml	18566	J43.1
Source:	UNIPROT ID:	
Armenian Hamster	Q02242	
Isotype:	Full Name:	
IgG	programmed cell death 1	

Applications

Tested Applications:
FC
Species Specificity:
Mouse

Background Information

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).


Storage

Storage:
Store at 2-8°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.09% sodium azide.

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ unstimulated or anti-CD3/CD28 stimulated (2 days) mouse splenocytes were surface stained with 0.5 ug Anti-Mouse PD-1/CD279 (65120-1-Ig, Clone: J43.1) and FITC Goat anti-hamster (Armenian) IgG Antibody at dilution 1:100, and 0.5 ug CoraLite® Plus 647 Anti-Mouse CD3 (17A2) (CL647-65077, Clone: 17A2). Cells were not fixed.