

For Research Use Only

# Anti-Mouse CD134/OX40 (OX-86)

Catalog Number: 65136-1-Ig



## Basic Information

Catalog Number:

65136-1-Ig

Size:

100ug , 500 µg/ml

Source:

Rat

Isotype:

IgG1, kappa

GenBank Accession Number:

BC065782

GeneID (NCBI):

22163

Full Name:

tumor necrosis factor receptor  
superfamily, member 4

Purification Method:

Affinity purification

CloneNo.:

OX-86

Recommended Dilutions:

FC: 1 ug per 10<sup>6</sup> cells in 100 µl  
suspension

## Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC : Con-A stimulated mouse splenocytes,

## Background Information

CD134, also known as OX40 and TNFRSF4, is a member of the TNFR-superfamily of receptors (PMID: 2828930; 9766631). It is a type I transmembrane protein predominantly expressed on activated T cells which include CD4 and CD8 T cells, Th2, Th1, and Th17 cells, as well as regulatory T cells (Tregs) (PMID: 20307208). CD134 is activated by its cognate ligand CD134L (OX40L) and functions as a T cell co-stimulatory molecule (PMID: 26215166). CD134-CD134L interactions have been proposed as a potential therapeutic target for treating autoimmune diseases, cancer and infectious disease (PMID: 26215166; 19426222).

## Storage

Storage:

Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10<sup>6</sup> Con-A stimulated BALB/c mouse splenocytes were surface stained with CoraLite® Plus 647 Anti-Mouse CD3 (CL647-65077, Clone: 17A2), 1 ug Anti-Mouse CD134 (65136-1-Ig, Clone: OX-86) or rat IgG1 isotype control, and FITC anti-rat IgG1 Antibody at dilution 1:100. Cells were not fixed.