For Research Use Only

Anti-Human CD73 (AD2) Catalog Number:65162-1-lg 2 Publications





Basic Information	Catalog Number: GenBank Accession Number: 65162-1-lg BC015940		cession Number:	Purification Method: Protein A purification	
	- 0		D1).	CloneNo.:	11
	Size: GeneID (NCBI): 100ug , 0.5 mg/ml 4907		51 <i>)</i> :	AD2	
	Source:	UNIPROT ID:			
	Mouse	P21589			
	Isotype:	otype: Full Name:			
	IgG1, kappa 5'-nucleotidase, ecto (CD73)				
		Calculated N 29 kDa, 63 kl			
Applications	Tested Applications:				
Applications	FC				
	Cited Applications:				
	FC				
	Species Specificity:				
	human				
	Cited Species: human				
	hemen				
Background Information	CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membrane- bound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).				
Notable Publications	Author	Pubmed ID	Journal		Application
	Junjie Liu	34707701	Exp Ther Med		FC
	Baihe Han	35968497	Dis Markers		FC
Storage	Storage: Store at 2-8°C. Stable for on Storage Buffer: PBS with 0.09% sodium azio				

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1X10^6 human PBMCs were surface co-stained with CL405 Anti-Human CD3 and 0.5 ug Anti-Human CD73 (65162-1-1g, Clone:AD2) and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 or Isotype Control. Cells were not fixed. Lymphocytes were gated.

1X10^6 human PBMCs were surface stained with 0.5 ug Anti-Human CD73 (65162-1-lg, Clone:AD2) and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red) or Isotype Control. Cells were not fixed. Lymphocytes were gated.