## For Research Use Only

## Anti-Rat CD8a (OX-8)

Catalog Number:65180-1-Ig



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

65180-1-lg

GeneID (NCBI):

Purified by protein-A affinity chromatography

Size:

24930

BC088126

CloneNo.:

**0X-8** 

100ug , 0.5 mg/ml Source:

UNIPROT ID:

Recommended Dilutions:

Mouse

P07725 Full Name:

FC: 0.5 ug per  $10^6$  cells in  $100 \mu l$ 

Isotype: IgG1, kappa

CD8a molecule

suspension

**Applications** 

**Tested Applications:** 

FC

**Positive Controls:** 

Species Specificity:

Rat

FC: Wistar rat splenocytes,

## **Background Information**

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR)  $\gamma\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

## Selected Validation Data



1X10^6 Wistar rat splenocytes were surface stained with 0.5 ug Anti-Rat CD8a (65180-1-lg, Clone:OX-8) and FITC anti-Mouse IgG1 Antibody at dilution 1:100, or 0.5 ug Control Antibody. Cells were not fixed.