

For Research Use Only

Anti-Human CD314/NKG2D (1D11)

Catalog Number: 65188-1-Ig



Basic Information

Catalog Number:

65188-1-Ig

Size:

100ug, 0.5 mg/ml

Source:

Mouse

Isotype:

IgG1, kappa

GenBank Accession Number:

BC039836

GeneID (NCBI):

22914

UNIPROT ID:

P26718

Full Name:

killer cell lectin-like receptor
subfamily K, member 1

Calculated MW:

25 kDa

Purification Method:

Purified by protein-A affinity
chromatography

CloneNo.:

1D11

Applications

Tested Applications:

FC

Species Specificity:

Human

Background Information

CD314, also known as NKG2D or Killer cell lectin-like receptor subfamily K member 1 (KLRL1), is a type II lectin-like transmembrane stimulatory receptor (PMID: 8436421). In humans, it is expressed on NK cells, gamma delta T cells, and CD8+ alpha beta T cells (PMID: 10426993). Various families of cell surface ligands have been identified, including the MIC A/MIC B and ULBP proteins (PMID: 12150888). CD314 is involved in both innate and adaptive immunities, and the NKG2D/NKG2DL pathway involves multiple effector cell types for controlling tumor progression (PMID: 31720075).

Storage

Storage:

Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human PBMCs were surface stained with 0.2 ug Anti-Human CD314/NKG2D (65188-1-Ig, Clone:1D11) or Mouse IgG1 Isotype Control and CoraLite®647-conjugated AffiniPure F(ab')₂ Fragment Goat Anti-Mouse IgG (H+L) (SA00014-8) at dilution 1:1000. Cells were then stained with CL488-conjugated CD8 (CL488-65135). Cells were not fixed. Lymphocytes were gated.



1x10⁶ human PBMCs were surface stained with 0.2 ug Anti-Human CD314/NKG2D (65188-1-Ig, Clone:1D11) or Mouse IgG1 Isotype Control and CoraLite®647-conjugated AffiniPure F(ab')₂ Fragment Goat Anti-Mouse IgG (H+L) (SA00014-8) at dilution 1:1000. Cells were not fixed. Lymphocytes were gated.