For Research Use Only

Anti-Human CD73 (AD2) Rabbit Recombinant Antibody



Catalog Number: 65500-1-RR

Basic Information

Catalog Number:

65500-1-RR

Size:

100ug , 0.5 mg/mL

Source:

Rabbit Isotype:

GenBank Accession Number:

GeneID (NCBI):

4907

BC015940

Full Name: 5'-nucleotidase, ecto (CD73)

Calculated MW:

29 kDa, 63 kDa

Purification Method: Affinity purification

CloneNo.: AD2

Applications

Tested Applications:

Species Specificity:

Human

Background Information

CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membranebound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).


Storage

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1X10^6 human PBMCs were surface stained with CL488 Anti-Human CD3 and 0.4 ug Anti-Human CD73 Rabbit Recombinant Antibody (65500-1-RR, Clone:AD2) or Rabbit IgG control PolyAb. Unconjugated antibodies were labeled with FlexAble CL750 Antibody Labeling Kit for Rabbit IgG (KFA004) prior to staining. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 0.4 ug Anti-Human CD73 Rabbit Recombinant Antibody (65500-1-RR, Clone:AD2) or Rabbit IgG control PolyAb. Unconjugated antibodies were labeled with FlexAble CL750 Antibody Labeling Kit for Rabbit IgG (KFA004) prior to staining. Cells were not fixed. Lymphocytes were gated.