

For Research Use Only

# Anti-Human CD11c (3.9) Mouse IgG1 Recombinant Antibody



Catalog Number: 65511-1-MR

## Basic Information

<b>Catalog Number:</b> 65511-1-MR	<b>GenBank Accession Number:</b> BC038237	<b>Purification Method:</b> Affinity purification
<b>Size:</b> 100ug, 0.5 mg/mL	<b>GeneID (NCBI):</b> 3687	<b>CloneNo.:</b> 3.9
<b>Source:</b> Mouse	<b>ENSEMBL Gene ID:</b> ENSG00000140678	
<b>Isotype:</b> IgG1	<b>Full Name:</b> integrin, alpha X (complement component 3 receptor 4 subunit)	
	<b>Calculated MW:</b> 1169 aa, 129 kDa	

## Applications

**Tested Applications:**  
FC

**Species Specificity:**  
Human

## Background Information

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated  $\alpha$  and  $\beta$  subunits (PMID: 9779984). CD11c, also known as integrin  $\alpha X$ , is a type I transmembrane glycoprotein present on a variety of cells, including monocytes/macrophages, granulocytes, a subset of B cells, NK cells and dendritic cells (PMID: 2897326; 1680915; 1694698; 17389580). As a result of its high level of expression on most dendritic cells, CD11c is typically considered to be a marker of conventional dendritic cells (PMID: 27119555). CD11c forms an  $\alpha/\beta$  heterodimer with CD18 (integrin  $\beta 2$ ). CD11c/CD18 acts a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis (PMID: 1671533).

## Storage


**Storage:**  
Store at 2-8°C. Stable for one year after shipment.

**Storage Buffer:**  
PBS with 0.09% sodium azide.


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>6</sup> human PBMCs were surface stained with 0.4 ug Anti-Human CD11c Mouse Recombinant Antibody (65511-1-MR, Clone:3.9) or Mouse IgG1 Isotype Control, and CoraLite®647-Conjugated AffiniPure Donkey Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were then stained with FITC-conjugated Anti-Human CD14. Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.



1x10<sup>6</sup> human PBMCs were surface stained with 0.4 ug Anti-Human CD11c Mouse Recombinant Antibody (65511-1-MR, Clone:3.9) or Mouse IgG1 Isotype Control, and CoraLite®647-Conjugated AffiniPure Donkey Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.