For Research Use Only

Anti-Human CD11c (3.9) Mouse IgG1 Recombinant Antibody

Catalog Number:65511-1-MR





Basic Information	Catalog Number: 65511-1-MR	GenBank Accession Number: BC038237	Purification Method: Affinity purification
	Size: 100ug , 0.5 mg/mL	GeneID (NCBI): 3687	CloneNo.: 3.9
	Source: Mouse	ENSEMBL Gene ID: ENSG00000140678	
	lsotype: IgG1	Full Name: integrin, alpha X (complement component 3 receptor 4 subunit)	
		Calculated MW: 1169 aa, 129 kDa	
Applications	Tested Applications: FC		
	Species Specificity: human		
Background Information	Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits (PMID: 9779984). CD11c, also known as integrin αX, is a type I transmembrane glycoprotein present on a variety of cells, including monocytes/macrophages, granulocytes, a subset of B cells, NK cells and dendritic cells (PMID: 2897326; 1680915; 1694698; 17389580). As a result of its high level of expression on most dendritic cells, CD11c is typically considered to be a marker of conventional dendritic cells (PMID: 27119555). CD11c forms an α/β heterodimer with CD18 (integrin β2). CD11c/CD18 acts a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis (PMID: 1671533).		
Storage	Storage: Store at 2-8°C. Stable for one Storage Buffer: PBS with 0.09% sodium azid		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10[^]6 human PBMCs were surface stained with 0.4 ug Anti-Human CD11c Mouse Recombinant Antibody (65511-1-MR, Clone:3.9) or Mouse IgG1 Isotype Control, and CoraLite®647-Conjugated AffiniPure Donkey Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were then stained with FITCconjugated Anti-Human CD14. Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.

1x10^6 human PBMCs were surface stained with 0.4 ug Anti-Human CD11c Mouse Recombinant Antibody (65511-1-MR, Clone:3.9) or Mouse IgG1 Isotype Control, and CoraLite®647-Conjugated AffiniPure Donkey Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.