For Research Use Only

Anti-Human CD107b / LAMP2 (H4B4) proteintech Rabbit IgG Recombinant Antibody

www.ptglab.com

Purification Method:

Affinity purification

CloneNo.:

H4B4

Catalog Number: 65520-1-RR

Basic Information

Catalog Number:

65520-1-RR

Size:

100ug , 0.5 mg/mL

Source: Rabbit

Isotype:

GenBank Accession Number:

BC002965 GeneID (NCBI):

3920

ENSEMBL Gene ID: ENSG0000005893

Full Name:

lysosomal-associated membrane

protein 2 Calculated MW: 45 kDa

Applications

Tested Applications:

FC (Intra)

Species Specificity:

Background Information

LAMP2 (CD107b) is a Lysosomal membrane glycoprotein. LAMP2 is extensively glycosylated with asparaginelinked oligosaccharides which protect it from intracellular proteolysis (PMID: 10521503). Although LAMP-2 is localized primarily in the endosome-lysosomal membrane of cells, it is also found on the plasma membrane under certain circumstances, e.g., after platelet activation, during granulocytic differentiation and activation, and in some tumor cells (PMID: 12221139). LAMP is involved in lysosomal stability and autophagy (PMID: 12221139). This glycoprotein provides selectins with carbohydrate ligands. LAMP2 may play a role in tumor cell metastasis (PMID 9426697).

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1x10^6 human PBMCs were intracellularly stained with 0.2 ug Anti-Human CD107b Rabbit Recombinant Antibody (65520-1-RR, Clone:H4B4) or Rabbit 1gG control Rabbit PolyAb, and CoraLite®647-Conjugated AffiniPure Donkey Anti-Rabbit 1gG(H+L) at dilution 1:1000. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Cells were treated with FC Receptor Block prior to staining.



1x10^6 human PBMCs were intracellularly stained with 0.2 ug Anti-Human CD107b Rabbit Recombinant Antibody (65520-1-RR, Clone:H4B4) or Rabbit 1gG control Rabbit PolyAb, and CoraLite®647-Conjugated AffiniPure Donkey Anti-Rabbit 1gG(H+L) at dilution 1:1000. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Cells were treated with FC Receptor Block prior to staining.