For Research Use Only

Anti-Human CD19 (4G7) Mouse IgG2a Recombinant Antibody

Catalog Number: 65562-1-MR



Purification Method:

Protein A purfication

CloneNo.:

4G7

Basic Information

Catalog Number:

65562-1-MR

Size:

100ug , 1 mg/ml

Source: Mouse Isotype: IgG2a

ENSG00000177455 UNIPROT ID: P15391 Full Name: CD19 molecule

> Calculated MW: 556 aa, 61 kDa

GenBank Accession Number:

BC006338

GeneID (NCBI):

ENSEMBL Gene ID:

Applications

Tested Applications:

Species Specificity:

human

Background Information

CD19 is a 95 kDa type I transmembrane glycoprotein belonging to the immunoglobulin superfamily (PMID: 2472450). It is expressed by B cells and follicular dendritic cells. CD19 is up-regulated at the step of B-lineage commitment during the differentiation of the hematopoietic stem cell, it remains on during subsequent stages of differentiation until finally down-regulated during terminal differentiation into plasma cells (PMID: 8528044). CD19 is involved in B cell development, activation and differentiation. It is the dominant component for the signaling complex on B cells that includes CD21 (CR2), CD81 (TAPA-1) and CD225 and acts as a critical co-receptor for BCR signal transduction (PMID: 23210908).


Storage

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD19 (4G7) Mouse IgG2a Recombinant Antibody (65562-1-MR, Clone: 4G7) or Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) and PE-conjugated Goat Anti-Mouse IgG. Cells were co-stained with APC Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (APC-65569, Clone: OKT3). Cells were not fixed. Lymphocytes were gated.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD19 (4G7) Mouse IgG2a Recombinant Antibody (65562-1-MR, Clone: 4G7) (red) or Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue), and PEconjugated Goat Anti-Mouse IgG. Cells were not fixed. Lymphocytes were gated.