For Research Use Only

Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody, PBS Only

Catalog Number:65569-1-PBS




Basic Information	Catalog Number: 65569-1-PBS	GenBank Accession Number: BC049847	Purification Method: Protein A purification
	Size: 1mg , 2mg/ml	Genel D (NCBI): 916	CloneNo.: OKT3
	Source: Mouse	ENSEMBL Gene ID: ENSG00000198851	
	Isotype: IgG2a	Full Name: CD3e molecule, epsilon (CD3-TCR complex)	
		Calculated MW: 207 aa, 23 kDa	
Applications	Tested Applications: FC		
	Species Specificity: human		
Background Information	CD3 is a multimeric protein associated with the T-cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction (PMID: 15885124). CD3 is composed of CD3 γ , δ , ϵ , and ζ chains (PMID: 1826255). It is expressed by thymocytes in a developmentally regulated manner, T cells, and some NK cells (PMID: 3289580). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).		
Storage	Storage: Store at -80°C. Stable for one year a Storage Buffer: PBS Only	fter shipment.	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^{^6} human PBMCs were surface stained with CoraLite[®] Plus 647 Anti-Human CD19, and 0.25 ug Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (65569-1-MR, Clone:OKT3) or 0.25 ug Mouse IgG2a Isotype Control (C1.18.4), (65208-1-Ig, Clone: C1.18.4), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. Lymphocytes were gated. This data was developed using the same antibody clone with 65569-1-PBS in a different storage buffer formulation.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (65569-1-MR, Clone: OKT3) (red) or 0.25 ug Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. Lymphocytes were gated. This data was developed using the same antibody clone with 65569-1-PBS in a different storage buffer formulation.