

For Research Use Only

Anti-Human CD64 (10.1) Mouse IgG2a Recombinant Antibody



Catalog Number: 65572-1-MR

Basic Information

Catalog Number: 65572-1-MR	GenBank Accession Number: BC032634	Purification Method: Protein A purification
Size: 100ug, 1 mg/ml	GeneID (NCBI): 2209	CloneNo.: 10.1
Source: Mouse	ENSEMBL Gene ID: ENSG00000150337	
Isotype: IgG2a	Full Name: Fc fragment of IgG, high affinity Ia, receptor (CD64)	
	Calculated MW: 374 aa, 43 kDa	

Applications

Tested Applications:
FC

Species Specificity:
Human

Background Information

Fcγ receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or inhibitory effects on cell functions after aggregation by complexed immunoglobulin G (IgG) (PMID: 17005690). CD64, also known as Fcγ R1A, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.

Storage


Storage:
Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.09% sodium azide.

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human PBMCs were surface stained with 0.5 ug Anti-Human CD64 (10.1) Mouse Recombinant Antibody (65572-1-MR, Clone:10.1) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1), and 0.5 ug Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4). Cells were then stained with APC Anti-Human CD11b (ICRF44) (APC-65116, Clone: ICRF44). Cells were not fixed. Monocytes were gated.